

# Interaction Between the *oxa1* and *rmp1* Genes Modulates Respiratory Complex Assembly and Life Span in *Podospora anserina*

Carole H. Sellem, Claire Lemaire, Séverine Lorin, Geneviève Dujardin  
and Annie Sainsard-Chanet<sup>1</sup>

Centre de Génétique Moléculaire UPR 2167, Associated with the University of Paris-Sud 11 CNRS, 91190 Gif-sur-Yvette, France

Manuscript received July 22, 2004  
Accepted for publication November 16, 2004

## ABSTRACT

A causal link between deficiency of the cytochrome respiratory pathway and life span was previously shown in the filamentous fungus *Podospora anserina*. To gain more insight into the relationship between mitochondrial function and life span, we have constructed a strain carrying a thermosensitive mutation of the gene *oxa1*. OXA1 is a membrane protein conserved from bacteria to human. The mitochondrial OXA1 protein is involved in the assembly/insertion of several respiratory complexes. We show here that *oxa1* is an essential gene in *P. anserina*. The *oxa1<sup>ts</sup>* mutant exhibits severe defects in the respiratory complexes I and IV, which are correlated with an increased life span, a strong induction of the alternative oxidase, and a reduction in ROS production. However, there is no causal link between alternative oxidase level and life span. We also show that in the *oxa1<sup>ts</sup>* mutant, the extent of the defects in complexes I and IV and the life-span increase depends on the essential gene *rmp1*. The RMP1 protein, whose function is still unknown, can be localized in the mitochondria and/or the cytosolic compartment, depending on the developmental stage. We propose that the RMP1 protein could be involved in the process of OXA1-dependent protein insertion.

**R**EGULATION of longevity and the processes that lead to age-related damages have been very actively studied in the past decade. Studies in model systems revealed numerous processes limiting longevity and aging. Among these processes, mitochondrial function seems to play a determinant role. The free radical theory, first proposed by HARMAN (1956), hypothesized that free radicals (reactive oxygen species, ROS) cause oxidative damage to proteins, lipids, and DNA, resulting in aging and death (SHIGENAGA *et al.* 1994). Mitochondria are the major source of ROS production and are also the major site of adenosine triphosphate (ATP) production. This dual role raises an endogenous challenge to the survival of the cell or organism. It is therefore not surprising that the genetic analysis of aging revealed a large number of genes essential for mitochondrial function. A causal link between respiration and life span was first established in the filamentous fungus *Podospora anserina* in which a loss of function of cytochrome *c* oxidase strikingly delays aging (DUFOUR *et al.* 2000). In the same way, several mutations that decrease the activity of the respiratory chain were shown to dramatically increase life span in *Caenorhabditis elegans* (FENG *et al.* 2001; LEE *et al.* 2003). Two recent studies in this organism also re-

vealed that disturbing the synthesis of subunits of the respiratory chain by RNA interference increases life span and that genes important for mitochondrial function stand out as the principal group of genes affecting longevity (DILLIN *et al.* 2002; LEE *et al.* 2003). Last but not least, a direct causal link between mitochondrial DNA mutations, respiratory chain dysfunction, and aging has just been demonstrated in mammals (TRIFUNOVIC *et al.* 2004). Altogether, these studies establish a strong link between low ATP content, low oxygen uptake, and increased longevity. However, the mechanisms by which mitochondrial function (ATP level, ROS level, complex metabolic changes, etc.) can regulate aging are not clearly understood.

*P. anserina* is a filamentous ascomycete that exhibits an unavoidable and specific arrest of vegetative growth called “senescence” (RIZET 1953). The senescent state is invariably associated with mitochondrial DNA (mtDNA) instability and accumulation of circular DNA molecules generated from the mitochondrial genome and called senDNAs (ALBERT and SELLEM 2002 and BELCOUR *et al.* 1999 for a review). As previously shown, inactivation of cytochrome *c* oxidase results in an increased mtDNA stability and longevity associated with decreased ROS production and deficits in energy-consuming processes such as fertility and growth rate (DUFOUR *et al.* 2000). Absence of the cytochrome part of the respiratory chain is not lethal in *P. anserina*, due to the activity of an alternative oxidase (AOX) known to branch off the cytochrome pathway at ubiquinone (VANLERBERGHE and

Sequence data from this article have been deposited with the EMBL/GenBank Data Libraries under accession no. AF382189.

<sup>1</sup>Corresponding author: Centre de Génétique Moléculaire UPR 2167, CNRS, 1 avenue de la terrasse, 91190 Gif-sur-Yvette, France.  
E-mail: sainsard@cgm.cnrs-gif.fr

MCINTOSH 1997; SLUSE and JARMUSZKIEWICZ 1998 for reviews). However, the exclusive use of the alternative pathway results in the loss of two of the three potential coupling sites and a decrease in ATP synthesis efficiency. Interestingly, the constitutive expression of the AOX pathway in long-lived respiratory mutants results in an increase of both ROS and ATP production (estimated from growth improvement and fertility restoration) correlated with a reduction of longevity (LORIN *et al.* 2001). All of these studies clearly point to mitochondrial energy as a key determinant of life span in *P. anserina* but do not allow us to specify what are the mitochondrial critical parameters that control life span.

In an attempt to gain more insight into the links between mitochondrial function and life span, we have studied the role of the nuclear *oxa1* gene. OXA1 is a mitochondrial membrane protein conserved from bacteria to chloroplasts and mitochondria (KUHN *et al.* 2003). It is a key component of the insertion machinery of membrane proteins (HERRMANN and NEUPERT 2003). In *Saccharomyces cerevisiae*, absence of Oxa1p activity is viable but mutants are unable to respire. They show severe defects in the assembly of cytochrome *c* oxidase (complex IV) and ATPase (complex V) associated with minor defects in the biogenesis of the *bc1* complex, complex III (BONNEFOY *et al.* 1994a; ALTAMURA *et al.* 1996; HAMEL *et al.* 1998). In *Schizosaccharomyces pombe*, the absence of the OXA1 protein results in a nonviable phenotype (BONNEFOY *et al.* 2000). These yeasts are devoid of complex I (proton-pumping NADH dehydrogenase). Recently, it has been shown in *Neurospora crassa* that OXA1 is also necessary for the biogenesis of complex I and that inactivation of the gene *oxa1* by the process of repeat-induced point mutation leads to lethality (NARGANG *et al.* 2002). The demonstration that at least some complex I subunits depend on OXA1 for their assembly/insertion extends the role of OXA1 to at least four of the five respiratory complexes.

In this study, we describe the properties of a *P. anserina* mutant that carries a deletion of the *oxa1* gene and an ectopic thermosensitive copy of the gene ( $\Delta$ *oxa1 oxa1<sup>ts</sup>*). Our results confirm that *oxa1* is an essential gene in obligate aerobic fungi, even in those that possess an alternative pathway, and that the OXA1 protein plays a key role in the biogenesis of complexes I and IV. Interestingly, we found a genetic interaction between the *oxa1* and *rmp1* genes. The *rmp1* gene exists under two natural alleles, *rmp1-1* and *rmp1-2*, that control the timing of death in strains carrying specific mutations in the nuclear gene *ASI*, which encodes a cytosolic ribosomal protein (DEQUARD-CHABLAT and SELLEM 1994). *rmp1* is an essential gene and the RMP1 protein, whose function has not been established, can be localized in the mitochondrial and/or the cytosolic compartment, depending on the developmental stage (CONTAMINE *et al.* 2004). We show here that the mutant form of OXA1

in the  $\Delta$ *oxa1 oxa1<sup>ts</sup>* strain grown at subrestrictive temperature leads to pleiotropic respiratory defects and to modification of life span. Interestingly, the extent of these alterations depends on the *rmp1* gene. The significance of the interaction between the *oxa1* and *rmp1* genes is discussed.

## MATERIALS AND METHODS

***P. anserina* strains, growth conditions, and transformation of protoplasts:** The genetics and biological properties of *P. anserina* were first described by RIZET and ENGELMANN (1949) and reviewed by Esser (ESSER 1974). Strains used in this study were derived from the *s* wild-type strain (RIZET 1952). The *leu1-1* mutant is auxotrophic for leucine. The *gpd-aox* strain contains an ectopic copy of the *aox* gene under the control of the strong constitutive *P. anserina gpd* promoter. The *aox::hygro* strain is inactivated for the endogenous *aox* gene (LORIN *et al.* 2001). The pHSS plasmid carries the hygromycin resistance gene and the *P. anserina rmp1-1* gene (CONTAMINE *et al.* 2004). All media, *i.e.*, cornmeal extract (MR), minimal synthetic (M2), and germination media, were as described (ESSER 1974). When necessary, hygromycin, phleomycin, and leucine were added to the M2 medium at a concentration of 75, 10, and 50  $\mu$ g/ml, respectively. Protoplast achievement and transformation experiments were conducted as described previously (BERGES and BARREAU 1989), except that protoplasts were obtained by incubation in 40 mg/ml glucanex (Laffort).

**Plasmids, bacterial, and yeast strains:** The plasmid used for the construction of the *P. anserina* cDNA libraries was the multicopy *S. cerevisiae* pFL61 expression vector, which carries the *URA3* gene as selectable marker in yeast (D'ENFERT *et al.* 1995). The plasmids used for constructions of the *P. anserina* strains bearing a deleted *oxa1* gene ( $\Delta$ *oxa1*) and/or an ectopic thermosensitive *oxa1* allele (*oxa1<sup>ts</sup>*) were pBC-hygro, which carries the bacterial *hph* gene conferring resistance to hygromycin as selectable marker (SILAR 1995), and pUL, which carries the wild-type *P. anserina leu1* gene (TURCQ 1989; DEBUCHY *et al.* 1993). The TOPO vector (Invitrogen, San Diego) was used to clone PCR amplification products. The pPable plasmid is a pUC vector containing the Tn5 bacterial bleomycin (*ble*) gene under the control of the *P. anserina gpd* promoter and the *S. cerevisiae cyc1* terminator (COPPIN and DEBUCHY 2000). The *ble* gene confers resistance to phleomycin. Plasmids were propagated in *Escherichia coli* XL1blue. Growth culture conditions and genetic and transformation methods for *S. cerevisiae* were as reported (BONNEFOY *et al.* 1994a).

**Construction of *P. anserina* cDNA libraries in the expression vector of yeast pFL61:** Total RNA (2 mg) was extracted from the *s* wild-type strain (*mat-*) and  $\sim$ 20  $\mu$ g of poly(A<sup>+</sup>) RNA was purified twice on oligo(dT)-cellulose columns (mRNA purification kit, Pharmacia). Anchored dT<sub>25</sub> primers were used to obtain 3.2  $\mu$ g double-strand DNA (cDNA kit, Amersham, Buckinghamshire, UK). Three cDNA libraries, corresponding to three ranges of molecular weight cDNA (0.2–1 kb, 1–2.5 kb, and >2.5 kb) were cloned using *Bst*X1 adaptators in the pFL61 vector between the 5' (promoter) and 3' (terminator) sequences of the *pgk1* gene as described previously (D'ENFERT *et al.* 1995). Sequencing of these cDNA libraries is currently in progress at Genoscope (Evry, France).

**Complementation of the yeast NBT2 mutant strain and isolation of the *oxa1* cDNA:** Since the pFL61 plasmid carries the *URA3* gene, the transformation events of the NBT2 strain—*oxa1::LEU2, leu2-3,112, ura3-1, his3-11,15, trp1-1, ade2-1, mat  $\alpha$*  (HAMEL *et al.* 1998)—can be easily selected by plating on a

fermentable medium (glucose) lacking uracil and containing histidine, tryptophan, and adenine. Among  $>10^5$  transformants screened, 50 and 5 transformants able to grow on nonfermentable carbon source (glycerol) were obtained from the 0.2- to 1-kb and the 1- to 2.5-kb cDNA libraries, respectively. Five of them were analyzed. They all contained a 1.2-kb insert whose sequence was identical and whose translated product is homologous to the OXA1 protein of several organisms.

**Cloning, sequencing, and localization of the *P. anserina* *oxa1* gene:** The *oxa1* cDNA used as a probe in Southern blot analysis revealed a 4.5-kb *NarI* DNA genomic fragment. A *NarI* mini-bank was thus constructed and screened with the same probe. The sequence of the 4478-bp cloned *NarI* fragment (Figure 1A) was determined and registered with the accession no. AF382189. A shorter (4-kb) *XhoI-NarI* fragment was subcloned in the pBC-hygro vector digested by *XhoI* and *ClaI*. The resulting pBCoxa1 plasmid was further used for various subclonings. The *oxa1* gene was located by hybridization on the separated chromosomes of *P. anserina* slotted on a nylon membrane (BARREAU *et al.* 2002) kindly provided by C. Barreau using the *oxa1* cDNA as a probe (chromoslot technique).

**Construction of strains containing an ectopic copy of the *oxa1* gene carrying a L269S substitution:** The construction of a transformation vector containing a thermosensitive *oxa1* allele was achieved as shown in Figure 1. First, a T-to-C transition in position 2651 was introduced in the open reading frame of the wild-type *oxa1* gene of pBCoxa1 to obtain a L269S substitution located in the loop between the transmembrane domains 2 and 3 of the OXA1 protein. This change corresponded to the L240S substitution introduced into the *S. cerevisiae* *oxa1* gene and resulted in the thermosensitive *pet*<sup>1402</sup> mutant (BAUER *et al.* 1994; MEYER *et al.* 1997a). This modification was achieved by direct mutagenesis in a two-step PCR amplification of the internal *NruI-PpuMI* DNA fragment (see legend of Figure 1B) and it was ascertained by sequencing. Second, the *NruI-PpuMI* of pBCoxa1 was excised and replaced by the modified *NruI-PpuMI* DNA to give pBCoxa1<sup>S</sup> (Figure 1B). Third, a 5.7-kb DNA fragment (*MluI-SpeI*) from pBCoxa1<sup>S</sup> containing the L269S allele was subcloned in the pUL vector digested by *AflIII* and *XbaI*. The resulting pULoxa1<sup>S</sup> vector (Figure 1C) was further used to transform the *leu1-1* strain to obtain ectopic integrations of the *oxa1*<sup>S</sup> gene (L269S *oxa1* allele). Five independent ectopic integration events of the plasmid were selected on minimal medium. These five primary transformants were crossed with the *leu1-1* strain to obtain prototroph monocaryotic transformed strains carrying the resident *oxa1*<sup>+</sup> gene and an ectopic L269S *oxa1* copy associated with both mating types. The *oxa1*<sup>+</sup> (*oxa1*<sup>S</sup>) resulting strain (the transgene in parentheses) had no observable phenotype and showed the same growth rate, mycelium aspect, and fertility as the wild-type strain at 27° as at 34°.

**Inactivation of the *oxa1* gene and construction of strains bearing the  $\Delta$ *oxa1* allele:** The construction of the pBC $\Delta$ oxa1::ble plasmid used to obtain a null allele of gene *oxa1* was achieved as follows: the internal *NruI-PpuMI* fragment of pBCoxa1 that contained most of the coding sequence of gene *oxa1* was excised and replaced by a 1.1-kb DNA fragment carrying the phleomycin resistance gene (*ble*). This construction gave the pBC $\Delta$ oxa1::ble vector (Figure 1D). A linear DNA fragment (*BspEI-BglIII*) of 3.7 kb issued from the pBC $\Delta$ oxa1::ble vector containing 1.4 kb upstream from the *NruI* site and 1.2 kb downstream from the *PpuMI* site was used to transform a *leu1-1 mat+* strain carrying an *oxa1*<sup>S</sup> ectopic copy. A total of 200 transformants, screened for phleomycin resistance, were tested by PCR analysis for the replacement of the wild-type *oxa1* resident gene by the inactivated  $\Delta$ *oxa1* allele. For this purpose, a couple of primers designed in the genomic sequence, upstream from

the *XhoI* restriction site (sense) and in the *ble* gene (antisense) were used (Figure 1, A and D). PCR amplifications were performed directly on crude extracts of lyophilized mycelia. Twelve transformants revealed a modified phenotype at 28° and 32° and 2 were slow growing only at 32°. One of the latter allowed the amplification of a fragment of the expected size.

**Isolation of mitochondria:** Mitochondria from cultures grown at 27° or 30° were isolated by differential centrifugation as described previously (SELLEM *et al.* 1990). Proteins were resuspended in 0.7 M sorbitol, 50 mM Tris-HCl, pH 7.5, and 0.2 mM EDTA containing protease inhibitor and further used for Western blot or BN-PAGE analysis. Bio-Rad (Richmond, CA) assay was used to determine protein concentration.

## RESULTS

**Characteristics of the *oxa1* sequence:** Cloning of the *P. anserina* *oxa1* cDNA by functional complementation in a *S. cerevisiae* *oxa1* null mutant revealed that the *P. anserina* OXA1 protein can rescue the yeast respiratory deficiency. The 426-amino-acid protein deduced from the *P. anserina* *oxa1* sequence shares 41, 21, and 22% identity with the OXA1 proteins from *N. crassa*, *S. cerevisiae*, and *Homo sapiens*, respectively. The same hydrophobic distribution of amino acids revealed the same characteristic structure with a conserved core region that probably also contains five transmembrane domains. A strong coiled-coil ( $\alpha$ -helix) structure, proposed to play a role in the recruitment of the mitochondrial translation machinery (SZYRACH *et al.* 2003), was predicted at the C-terminal domain of the *P. anserina* protein as in that of *N. crassa* and *S. cerevisiae*. The *oxa1* gene was cloned and modified as described in MATERIALS AND METHODS (see Figure 1). The phleomycin-resistant phenotype associated with the *oxa1* inactivation allowed us to genetically localize the *oxa1* gene far from its centromere (second-division segregation percentage of  $\sim$ 90). It was assigned to chromosome II by the chromoslot technique.

**The inactivation of *oxa1* is lethal in *P. anserina*:** A *mat+* strain bearing the  $\Delta$ *oxa1* allele was obtained as described in MATERIALS AND METHODS. The recipient strain used for the transformation-mediated gene replacement carried an ectopic copy of the *oxa1* gene with an L269S substitution (*oxa1*<sup>S</sup>). The primary transformant strain,  $\Delta$ *oxa1 mat+* (*oxa1*<sup>S</sup>), was crossed at 27° with a *leu1-1 mat-* strain. In the progeny, we looked for asci-containing ascospores bearing the  $\Delta$ *oxa1* allele (determined by resistance to phleomycin) without the *oxa1*<sup>S</sup> ectopic copy (determined by auxotrophy for leucine). Among  $\sim$ 50 asci analyzed (each containing two monocaryotic spores), 8 asci revealed nonambiguously that homocaryotic spore resistant for phleomycin and auxotroph for leucine were unable to germinate. The fact that all germinating spores of the phleomycin-resistant phenotype are also leucine prototrophs indicates that they can survive only when the *oxa1*<sup>S</sup> transgene covers for the deleted allele and allows us to conclude that the inactivation of the *oxa1* gene is lethal for *P. anserina*.




FIGURE 1.—Cloning and inactivation of the *oxa1* gene. (A) Partial restriction map of the *P. anserina* wild-type *oxa1* gene. Nomenclature is according to the submitted sequence (no. AF 382189). The arrows represent the oligonucleotides used for further PCR experiments with their 5′–3′ orientation. (B) Partial restriction map of the *pBCoxa1<sup>ts</sup>* vector. The *XhoI*–*NarI* fragment of the *P. anserina* wild-type *oxa1* gene was cloned in the *pBC*–*hygro* vector (*pBCoxa1*). An *NruI*–*PpuMI* modified fragment obtained in a two-step PCR amplification replaced the *NruI*–*PpuMI* wild-type fragment. The first step was the amplification of two fragments using two couples of primers (*NruI*/C2 and C1/*PpuMI*), including two complementary oligonucleotides (C1 and C2) designed with the desired T-to-C substitution. In the second step, the *NruI*/*PpuMI* fragment was amplified using the two preceding amplified fragments as matrices. (C) Partial restriction map of the *pULoxa1<sup>ts</sup>* vector. The *MluI*–*SpeI* fragment of *pBCoxa1<sup>ts</sup>* was cloned in a *pUL* vector carrying the *leu1* gene. (D) Partial restriction map of the *pBCΔoxa1:ble* vector. The *NruI*–*PpuMI* wild-type fragment of *pBCoxa1* was replaced by an *Ecl136II*–*HindIII* DNA fragment carrying the *ble* gene using a *HindIII*/*PpuMI* linker. The shaded boxes correspond to the *oxa1* coding sequence. Dashed and dotted lines correspond to plasmid sequences. Ap, Cm, and *hygro* are bacterial genes conferring, respectively, the ampicillin, chloramphenicol, and hygromycin resistance to the *pUC*- and *pBC*–*hygro*-derived vectors.

**An L269S substitution in the gene *oxa1* confers a thermo-sensitive phenotype with pleiotropic effects and modified longevity at subrestrictive temperature:** The  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) mutant was grown at several temperatures ranging from 27° to 34°, and the life span, growth rate, and female fertility were analyzed. As shown in Table 1, this mutant cannot grow at 34°, and 31° was the highest temperature compatible with growth indicating that, as in *S. cerevisiae*, the L269S substitution confers a thermosensitive phenotype. Life-span determination revealed no effect of the mutation at 27° whereas at the subrestrictive temperature of 30°, the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>-</sup>* cultures displayed a reduced life span and the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>+</sup>* cultures an increased one (at least 250% of the wild type). The life span of  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>+</sup>* cultures appeared extremely heterogenous (see the standard deviation in Table 1) and the arrests of growth did not correspond to death of the mycelium, which always resumed growth after transfer to 27°. In contrast, the arrests of growth

at 27° and 30° of the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>-</sup>* and at 27° of the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>+</sup>* strains were due to death and the apical cells were unable to grow when transferred at 27°. The growth rate was almost unaffected at 27° in  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) strains, irrespective of the mating type, and only slightly reduced for the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>+</sup>* strain grown at 30°. In contrast, female fertility appeared very reduced even at 27° and this defect was much more marked at 30° where female fertility of the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) strains was <10% of that of the wild-type strain.

Growth arrest in the wild-type strain is due to senescence and systematically correlated with the accumulation of circular, tandemly arranged mtDNA called senDNAs (BELCOUR *et al.* 1999). Whereas senDNAs were revealed in  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>-</sup>* and *mat<sup>+</sup>* cultures dying at 27° and in  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>-</sup>* cultures dying at 30°, arrest of growth of the  $\Delta oxa1$  (*oxa1<sup>ts</sup>*) *mat<sup>+</sup>* cultures at 30° was not accompanied with such accumulation of senDNAs (data not shown). This result favors the idea that the

TABLE 1

Growth characteristics of the strains at permissive (27°), subrestrictive (30°), and restrictive (34°) temperatures

Strain	Lifespan (cm)			Growth rate (cm/day)			Female fertility	
	27°	30°	34°	27°	30°	34°	27°	30°
Wild-type <i>mat</i> <sup>-</sup>	8.1 ± 1.3 (6)	9.1 ± 1.7 (9)	7.0 ± 1.5 (3)	0.82 ± 0.05	0.81 ± 0.09	0.90 ± 0.11	100	100
Wild-type <i>mat</i> <sup>+</sup>	9.1 ± 3.7 (7)	10.8 ± 2.5 (6)	5.1 ± 2.6 (3)	0.80 ± 0.06	0.80 ± 0.03	0.89 ± 0.22	100	100
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) <i>mat</i> <sup>-</sup>	5.8 ± 2.2 (13)	5.1 ± 1.8 (15)	–	0.80 ± 0.10	0.75 ± 0.05	–	38	8
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) <i>mat</i> <sup>+</sup>	9.3 ± 3.8 (13)	28.1 ± 17.0 (17)	–	0.77 ± 0.08	0.70 ± 0.09	–	27	5
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) ( <i>rmp1-1</i> ) <i>mat</i> <sup>-</sup>	6.1 ± 0.5 (2)	5.2 ± 0.5 (2)	–	–	–	–	–	–
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) ( <i>rmp1-1</i> ) <i>mat</i> <sup>+</sup>	4.2 ± 2.5 (4)	4.6 ± 0.5 (5)	–	–	–	–	–	–
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) ( <i>gpd-aox</i> ) <i>mat</i> <sup>-</sup>	10.5 ± 3.4 (13)	10.6 ± 3.3 (5)	–	–	–	–	–	–
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) ( <i>gpd-aox</i> ) <i>mat</i> <sup>+</sup>	11.6 ± 3.5 (14)	43.2 ± 23.6 (10)	–	–	–	–	–	–
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) ( $\Delta$ <i>aox</i> ) <i>mat</i> <sup>-</sup>	7.0 ± 0.5 (3)	7.0 ± 0.7 (3)	–	–	–	–	–	–
$\Delta$ <i>oxa1</i> ( <i>oxa1</i> <sup>ts</sup> ) ( $\Delta$ <i>aox</i> ) <i>mat</i> <sup>+</sup>	8.7 ± 1.2 (3)	28.1 ± 20.7 (14)	–	–	–	–	–	–

To determine life span, cultures were grown on race tubes containing minimal synthetic M2 medium with twice the usual amount of agar at various temperatures ranging from 27° to 34°. Life spans ( $\pm$  standard deviation) are mean values determined in centimeters of growth initiated from the ascospore to the arrested edge of 2–17 parallel cultures of 1–10 different ascospores of the same genotype (the total number of cultures is given in parentheses). –, absence of growth. For each strain, growth rate was estimated on 5–15 subcultures over 5 days of growth and expressed in centimeters of growth per day. Female fertility was estimated by the number of mature perithecia that developed when the strains were fertilized with a suspension of wild-type conidia of the opposite mating type and is expressed as the percentage of the fertility of the wild-type strain of the same mating type at the same temperature. Transgenes are indicated in parentheses. The resident *rmp1* allele is not indicated in the table: all the *mat*<sup>-</sup> strains are *rmp1-1*; all the *mat*<sup>+</sup> strains are *rmp1-2*.

arrest of growth in the cultures with increased life span ( $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) *mat*<sup>+</sup> at 30°) is not the result of a senescence process.

**Gene *oxa1* interacts with the nuclear mitochondrial gene *rmp1*:** In *P. anserina*, a gene closely linked to the *mat* locus (<0.5 cM) has been identified (CONTAMINE *et al.* 1996, 2004). This gene, *rmp1*, exists under two natural alleles in strain *s*: *rmp1-1* and *rmp1-2*, linked, respectively, to the *mat*<sup>-</sup> and *mat*<sup>+</sup> loci. Gene *rmp1* controls the timing of premature death, a syndrome characterized by the accumulation of specific mitochondrial deletions (BELCOUR *et al.* 1991). This syndrome is observed only in strains carrying some mutations in the *ASI* gene, such as *ASI-4*, which alter a cytosolic ribosomal protein (DEQUARD-CHABLAT *et al.* 1986). Both *ASI-4 rmp1-1* and *ASI-4 rmp1-2* strains accumulate specific mitochondrial deletions at time of death, but they strongly differ in their longevity, which is ~2 and 80 cm, respectively (CONTAMINE *et al.* 1996). Gene *rmp1* has been shown to be an essential gene encoding a protein that can be localized in the mitochondrial and/or the cytosolic compartments (CONTAMINE *et al.* 2004). So, we asked whether the differences observed between  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) *mat*<sup>+</sup> and *mat*<sup>-</sup> strains at 30° revealed an interac-

tion between the *oxa1* gene and the *mat* locus or between the *oxa1* gene and the *rmp1* gene. To address this question, we took advantage of the cloning of the *rmp1-1* allele on an hygromycin transformation vector and of the dominance of this allele (CONTAMINE *et al.* 2004) to introduce *rmp1-1* by transformation into a  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) *rmp1-2 mat*<sup>+</sup> strain. Secondary  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) (*rmp1-1*) *rmp1-2 mat*<sup>+</sup> transformants were obtained and analyzed. Interestingly, as shown in Table 1, this analysis revealed that, when introduced into a  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) *rmp1-2 mat*<sup>+</sup> strain, *rmp1-1* leads to a reduced life span, not different from that of  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) *rmp1-1 mat*<sup>-</sup>, and to a restoration of accumulation of senDNA at the time of death (not shown). These data clearly demonstrate that the timing of death and the instability of the mtDNA in the  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) strains grown at subrestrictive temperature depend on the *rmp1* gene and not on the *mat* locus.

**The activity of the respiratory complexes I and IV is differentially impaired in  $\Delta$ *oxa1* (*oxa1*<sup>ts</sup>) *rmp1-1* and *rmp1-2* strains:** As demonstrated in several organisms, the OXA1 protein is required for the biogenesis of several respiratory complexes, namely complexes IV and V (BONNEFOY *et al.* 1994a; ALTAMURA *et al.* 1996) and complex I (NARGANG *et al.* 2002). The effect of the *oxa1*<sup>ts</sup>




FIGURE 2.—In-gel detection of complex I, IV, and V activities in various strains grown at 30°. Sample preparation and blue-native polyacrylamide gel electrophoresis (BN-PAGE) were carried out as described (SCHAGGER and VON JAGOW 1991). Mitochondria from *P. anserina* were rendered soluble in the presence of 2% (w/v) *N*-dodecyl maltoside. In-gel activity assays of complexes I, IV, and V were performed as described (NIJTMANS *et al.* 2002). Genotypes of the strains are detailed below the gel. The *oxa1* gene is either present (+) or inactivated (Δ). + or - indicates the presence or the absence of the *(oxa1<sup>ts</sup>)* and *(rmp1-1)* transgenes. The two *mat* alleles, *mat+* and *mat-*, are indicated by + and -, respectively. The specific activity of cytochrome *c* oxidase (COX) is expressed as the percentage of the wild type (wild type is 856 ± 54 nmol of oxidized cytochrome *c* per minute per milligram of mitochondrial protein). The wild-type strain (lane 1) gave the same results irrespective of *mat* and *rmp1* haplotype.

mutation on the activity of these different complexes was investigated in *P. anserina*  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) strains grown at 30°. Blue-native polyacrylamide gel electrophoresis (BN-PAGE), which allows a one-step separation of the different complexes in a gel activity assay, was performed (Figure 2). As expected from the corresponding *S. cerevisiae* *pet*<sup>1402</sup> mutant (MEYER *et al.* 1997a), complex V did not seem to be significantly affected in *P. anserina*  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) strains despite the information at or near the *mat* haplotype (lanes 2 and 3). In the same extracts a very clear difference was observed between  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-1* *mat-* and  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-2* *mat+* for complex I assembly/activity, which was only slightly affected, if at all, in the *rmp1-1* *mat-* strain (lane 2) or, in contrast, strongly reduced in the *rmp1-2* *mat+* strain (lane 3). In the same way, it seemed that complex IV assembly/activity was lower in  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-2* *mat+* than in  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-1* *mat-* strains. To verify this difference, specific activities of complex IV were determined on isolated mitochondria. They were evaluated to correspond, respectively, to ~20% for *rmp1-2* *mat+* and 70% for *rmp1-1* *mat-* of the wild-type strain. Interestingly, the analysis of the  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-2* *mat+* strain containing the *rmp1-1* transgene (lane 4) revealed a restoration of both complexes I and IV, indicating that




FIGURE 3.—Immunodetection of AOX by Western blot analysis. Twenty micrograms of mitochondrial proteins extracted from cultures grown at 30° were fractionated by SDS-PAGE and transferred to nitrocellulose membranes. Membranes were probed first with an anti-AOX mouse monoclonal antibody generated against the AOX of *Sauromatum guttatum* (ELTHON *et al.* 1989). Blots were then reprobed with an anti-βATPase rabbit antibody as a standardization control. The bound antibodies were detected using an enhanced chemiluminescence detection system (Pierce, Rockford, IL). For genotypes see legend of Figure 2. The percentage of respiration sensitive to cyanide (KCN, 1 mM) and SHAM (1 mg/ml) reflects the engagement of electrons in the cytochrome and alternative pathway, respectively. Oxygen uptake measurements were performed on protoplast suspensions (10<sup>8</sup> protoplasts/ml in 0.6 M sorbitol) in an oxytherm chamber with a Clark-type O<sub>2</sub> electrode (Hansatech). Wild-type strains used as controls (lane 1) gave the same results irrespective of their *mat* and *rmp1* haplotype.

in *P. anserina* the biogenesis of these complexes depends on both OXA1 and RMP1 proteins.

**The alternative pathway is differentially induced in  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-1* and *rmp1-2* strains:** It is well established that absence of the Oxa1p function leads to a drastic reduction in complex IV activity in *S. cerevisiae* and *S. pombe* and that decreased OXA1 level leads to reduced content in some subunits of this complex in *N. crassa* (NARGANG *et al.* 2002). The results presented here demonstrate that in *P. anserina* a decreased activity of complex IV is also observed in  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) strains. In this organism, impairment in complex IV is known to cause the induction of the alternative pathway, whose expression is practically null in wild type (DUFOUR *et al.* 2000; BORGHOUTS *et al.* 2001; LORIN *et al.* 2001).

To investigate whether the alternative oxidase is induced in the  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) strains, immunodetection of the AOX protein by Western blot analyses was performed. The analysis revealed that when mitochondria were extracted from cultures grown at 27°, no AOX could be detected in the wild-type or the mutant  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *mat+* *rmp1-2* and *mat-* *rmp1-1* strains. In contrast, as shown in Figure 3, when mitochondria were

extracted from cultures grown at 30°, there was no detectable protein in the wild type (lane 1) whereas a very low or a high amount of the AOX protein was present in  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-1$   $mat^-$  (lane 4) and  $rmp1-2$   $mat^+$  (lane 2), respectively. The level of induction of AOX in the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$  strains at 30° was estimated to be equivalent to that observed in strains containing the *aox* gene under the transcriptional control of the strong constitutive *gpd* promoter (lane 5). It is worth noting that, as for the other parameters tested, introduction of *rmp1-1* in  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  restored a  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-1$   $mat^-$  phenotype and led to a spectacular decrease in the AOX level (lane 3).

To ensure that the alternative pathway induced in  $\Delta oxa1$  ( $oxa1^{ts}$ ) strains is functional, the relative extent to which specific inhibitors of the alternative (salicylhydroxamic acid, SHAM) or cytochromic (potassium cyanide, KCN) pathways inhibit oxygen uptake was measured. As shown at the bottom of Figure 3, respiration of the wild type grown at 30° was ~80% and 20% sensitive to KCN and SHAM, respectively. In contrast, respiration of the  $\Delta oxa1$  ( $oxa1^{ts}$ ) strains was more sensitive to SHAM and less sensitive to KCN and this particularity was much more marked for the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  strains whose respiration was ~30% and 70% sensitive to KCN and SHAM, respectively. This clearly indicated that, in the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  strains grown at subrestrictive temperature, the alternative pathway was the most important way of transferring the electrons to oxygen. These results are in agreement with the Western blot analysis and it can be concluded that there is either a very strong or a faint induction of the alternative pathway in the  $\Delta oxa1$  ( $oxa1^{ts}$ ) strains grown at subrestrictive temperature according to the *rmp1* haplotype.

**The level of expression of the alternative oxidase does not causally control the phenotypic differences between  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-1$  and  $rmp1-2$ :** Relationship between life span and the alternative oxidase level in cytochrome-deficient *P. anserina* mutants is far from clear. So we decided to test whether or not the differences in life span observed between the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-1$  and  $rmp1-2$  strains are causally related to the level of expression of the AOX protein. First, to address this question,  $\Delta oxa1$  ( $oxa1^{ts}$ ) strains carrying the *aox* inactivated gene (*aox::hygro*) or an ectopic copy of the *aox* gene under the control of the strong constitutive *gpd* promoter (*gpd-aox*) were constructed by crossing  $\Delta oxa1$  ( $oxa1^{ts}$ ) with the *aox::hygro* ( $\Delta aox$ ) and the *gpd-aox* strain (LORIN *et al.* 2001), respectively. The analysis of these crosses first revealed that  $\Delta oxa1$  (*gpd-aox*) spores (determined by phleomycin and hygromycin resistance) did not germinate and that  $\Delta oxa1$  ( $oxa1^{ts}$ ) (*gpd-aox*)  $rmp1-2$   $mat^+$  and  $rmp1-1$   $mat^-$  strains did not grow beyond 31°, indicating that the constitutive expression of the *gpd-aox* gene does not rescue the lethal phenotype conferred by the *oxa1* gene inactivation. Second, the determination of the life span of the strains carrying the *gpd-aox* transgene re-

vealed, as shown in Table 1, that although the constitutive expression of AOX seemed to slightly improve life span in both  $\Delta oxa1$  ( $oxa1^{ts}$ ) (*gpd-aox*)  $rmp1-1$   $mat^-$  and  $rmp1-2$   $mat^+$  strains, it did not confer to  $\Delta oxa1$  ( $oxa1^{ts}$ ) (*gpd-aox*)  $rmp1-1$   $mat^-$  strains the long life span of  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  strains. We have verified that the *gpd-aox* transgene was efficiently expressed in the  $\Delta oxa1$  ( $oxa1^{ts}$ ) (*gpd-aox*)  $rmp1-1$   $mat^-$  strain, as shown in Figure 3 (lane 5). The functionality of the *gpd-aox* transgene is attested to by the increased sensitivity to SHAM (70%). Third, as shown in Table 1, the inactivation of the *aox* gene in  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$   $\Delta aox$  strains did not significantly reduce life span. Altogether, these data indicate that the different level of expression of the AOX protein between  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  and  $rmp1-1$   $mat^-$  strains at 30° is not the causal factor that accounts for the difference in longevity between the two strains.

**ROS production is lowered in  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$  strains:** In *P. anserina*, absence of complex IV activity results in an increased life span and a decreased production of reactive oxygen species (DUF0UR *et al.* 2000). To test whether there is a correlation between life span and production of ROS in the  $\Delta oxa1$  ( $oxa1^{ts}$ ) strains, ROS production was measured by cytofluorometry on protoplasts harvested from cultures grown at 30°. Figure 4A shows the results of three independent experiments indicating that the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  strain produced 50–60% less ROS than the wild type did while the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-1$   $mat^-$  strain produced as much or more. These data are in good agreement with the histochemical superoxide radical detection shown in Figure 4B. In this test, the relative level of produced superoxide radicals was visualized by the relative ability of mycelia to reduce yellow water-soluble nitro-blue tetrazolium (NBT) to blue water-insoluble formazan (MUNKRES 1990). The amount of secreted superoxide radicals seems to be equivalent in wild-type and  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  and  $rmp1-1$   $mat^-$  strains grown at 27°. In contrast, it is dramatically reduced in the  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-2$   $mat^+$  strain grown at 30° compared to the wild-type and  $\Delta oxa1$  ( $oxa1^{ts}$ )  $rmp1-1$   $mat^-$  strains grown under the same conditions. It appears to be equivalent to that observed in the complex IV-deficient long-lived mutant *cox5::ble*, which strengthens the idea of a strong link between life span and ROS production in *P. anserina*.

## DISCUSSION

In this study, we have identified the *oxa1* gene of *P. anserina*. It is functionally homologous to the *S. cerevisiae* *oxa1* gene and is essential for viability in *P. anserina*. A strain carrying a thermosensitive copy of the gene ( $oxa1^{ts}$ ), grown at subrestrictive temperature, exhibits severe defects in the respiratory complexes I and IV. One of the most striking results obtained in this study is the




FIGURE 4.—ROS measurements. (A) ROS production estimated as the kinetics of dichlorofluorescein (DCF) production resulting from the oxidation of the diacetate form ( $H_2DCF-DA$ ) as described previously (DUFOUR *et al.* 2000). For each strain grown at 30°,  $\sim 10^7$  protoplasts were incubated with  $H_2DCF-DA$  (80  $\mu M$  in 0.6 M sorbitol/10 mM Tris-HCl pH 7.5) at 30° and ROS production was analyzed by measuring, every 10 min, the apparition of the fluorescent DCF in a XL3C flow cytometer (Coulter, France). (B) Histochemical detection of superoxide radicals on mycelium (MUNKRES 1990). Cultures were grown at 27° or 30° on M2 medium until a diameter of 3 cm was reached: the petri dishes were flooded with 3–5 ml of staining solution containing 2.5 mM NBT diluted in 5 mM 3-(*N*-morpholino) propane sulfonate-NaOH (MOPS) buffer, pH 7.6. After an incubation of 60 min at 30°, the solution was discarded and plates kept in the dark. A blue, water-insoluble formazan precipitate, obtained from the reduction of the water-soluble NBT, visualized superoxide radicals. rmp1-1, rmp1-2,  $\Delta cox5$ , and wt, respectively, correspond to  $\Delta oxa1(oxa1^{ts}) rmp1-1 mat^-$ ,  $\Delta oxa1(oxa1^{ts}) rmp1-2 mat^+$ ,  $\Delta cox5 mat^+$ , and the wild-type  $mat^+$  strains. These last two strains gave the same results irrespective of their *mat* and *rmp1* haplotype.

existence of an interaction between genes *oxa1* and *rmp1*. The effects of complex I and IV activity impairment are an increased life span, a strong induction of the alternative oxidase, and a decrease in ROS production. Results presented here show that there is no causal link between the increase of life span and the amount of alternative oxidase.

**The *P. anserina* OXA1 protein is functionally homologous to the *S. cerevisiae* protein and is essential for viability:** The *P. anserina* OXA1 protein is able to complement the respiratory defect of a *S. cerevisiae* null mutant, as was previously shown for the OXA1 proteins from *S. pombe* (BONNEFOY *et al.* 2000), *N. crassa* (NARGANG *et al.* 2002), *Arabidopsis thaliana* (HAMEL *et al.* 1997), and *H. sapiens* (BONNEFOY *et al.* 1994b), demonstrating that all these proteins are truly functional homologs. In *S. cerevisiae*, a single transition mutation that results in an L240S substitution in a hydrophilic region between the second and third transmembrane domains leads to a respiratory thermosensitive phenotype (BAUER *et al.* 1994; MEYER *et al.* 1997b). The Leu240 residue is conserved in the different OXA1 homologs, ranging from bacteria to human (MEYER *et al.* 1997b). We show here that a change of the corresponding leucine to serine (L269S) in the *P. anserina* protein also leads to a thermosensitive phenotype, supporting the view that this conserved residue is essential for the function of OXA1 proteins.

The absence of Oxa1p in *S. cerevisiae* and that of the Sp2 ortholog in *S. pombe* lead to a complete loss of respiration. However, the knockout of the *oxa1* gene in *S. cerevisiae* is viable (BONNEFOY *et al.* 1994a), whereas the knockout of the two *S. pombe* orthologs, *Sp1* and *Sp2*, is nonviable (BONNEFOY *et al.* 2000). It was proposed that this lethality is linked to the small negative phenomenon and that the loss of Oxa1p function leads to severe defects in the F1-ATPase, resulting in the incapacity to ensure a basal level of membrane potential (BONNEFOY *et al.* 2000). The OXA1 function is of course expected to be essential for viability in strictly aerobic organisms, in which all the electron flow is transported through the cytochrome pathway. In organisms possessing an alternative respiration system branched to the cytochrome pathway, such as the filamentous fungi, the question of the OXA1 function requirement deserved to be raised. However, in both *N. crassa* and *P. anserina* the absence of OXA1 was shown to be nonviable. The lethality of  $\Delta oxa1$  mutants cannot be explained by defects in complex IV biogenesis since loss of function of cytochrome *c* oxidase is viable in *P. anserina*. It can therefore be concluded that it is due to complex V defects as hypothesized for *S. pombe* and/or to complex I defects. This complex, absent in *S. cerevisiae* and *S. pombe*, is situated upstream of the branch point of the alternative oxidase and therefore cannot be compensated by it. In *N. crassa*, depletion of OXA1 leads to



reduced levels of the 24- and 29.9-kD subunits of complex I revealed in Western blot (NARGANG *et al.* 2002). We show here that in a *rmp1-2* context, a compromised OXA1 activity reduces complex I activity dramatically and, to a lesser extent, complex IV activity, whereas it has practically no effect on complex V activity.

**Genetic interaction between the *oxa1* and *rmp1* genes:**

A remarkable and exciting result provided by this work is the occurrence of a genetic interaction between the *oxa1* and *rmp1* genes. The two natural alleles *rmp1-1* and *rmp1-2* differ by three substitutions but the life-span differences observed between *AS1-4 rmp1-1* and *AS1-4 rmp1-2* have been attributed to a premature stop, which yields a protein lacking its last 19 amino acids in the *rmp1-2* context. Several lines of evidence support the hypothesis that *rmp1-1* is the fully functional allele (CONTAMINE *et al.* 2004). Interestingly, the defects in the biogenesis of complexes I and IV in  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) grown at 30° are much more pronounced in a *rmp1-2* than in a *rmp1-1* context, suggesting that the mutant form of OXA1 displays dependency on the RMP1 functional protein to play its role in the biogenesis of the respiratory complexes. The synergic relationship between the mutant OXA1 and the supposed not fully functional form of RMP1 suggests that the two encoded proteins are involved in the same process. The function of RMP1 is unknown and putative homologs have been found only in multicellular ascomycetes (CONTAMINE *et al.* 2004). However, the authors hypothesize that the RMP1 function could be widely distributed and could have evolved so rapidly that recognition of its homologs in distant species would be impaired. These authors have reported database searches indicating that the *Histoplasma capsulatum* homolog to RMP1 presents a putative homolog in *S. pombe*, which itself shows a weak similarity to the *S. cerevisiae* Sls1p protein. Sls1p is a mitochondrial membrane protein required for respiration (ROUILLARD *et al.* 1996). Its precise function remains to be elucidated. Recent data indicate that Sls1p is involved in mitochondrial translation regulation in cooperation with Nam1p and could serve to coordinate transcription and translation of mtDNA encoded products (BRYAN *et al.* 2002; RODEHEFFER and SHADEL 2003). It was also recently reported that Oxa1p may serve as a ribosome-binding site that allows coupling of the synthesis and membrane integration of mtDNA encoded proteins (JIA *et al.* 2003; SZYRACH *et al.* 2003). The two proteins could therefore interact at least during some cellular stages. Furthermore, on the basis of the fact that overexpression of *SLS1* does not rescue the translation defect of mutants of the amino terminal domain of the mtRNA polymerase but increases the steady-state level of some mitochondrial encoded proteins, it has been proposed that Sls1p could be involved in the assembly of the oxidative phosphorylation complexes in a manner independent of its effects on translation (RODEHEFFER and SHADEL 2003). Regardless the significance of these observations, the genetic interaction between the *oxa1*

and *rmp1* genes revealed in this study points to a role for the RMP1 protein in the process of OXA1-dependent protein insertion and provides a new framework for further studies.

**Impairment of complex I and IV activities leads to increased life span and strong induction of the alternative oxidase but these two parameters are not causally linked:**

This study confirms the influence of the mitochondrial activity in the control of life span and is consistent with previous work using respiratory-deficient mutants (DUFOUR *et al.* 2000; LORIN *et al.* 2001; STUMPFERL *et al.* 2004). In the  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-1* strains grown at subrestrictive temperature, the defects in complexes I and IV are nonexistent or modest, and life span is slightly reduced. In the  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-2* strains, in contrast, complexes I and IV appear strongly affected, and life span is considerably extended. Defects in complexes I and IV activities in *rmp1-2* strains are paralleled with a strong induction of the alternative oxidase, a shift of the partitioning of the electrons through this oxidase, leading to a highly SHAM-sensitive respiration and to a decrease in ROS production (DUFOUR *et al.* 2000; LORIN *et al.* 2001; STUMPFERL *et al.* 2004). If there is no doubt that close links occur between life span, AOX level, and ROS production in respiratory-deficient mutants of *P. anserina*, the nature of these links is still obscure. A correlation between increased life span and induced *aox* expression has been recently reported (BORGHOUTS *et al.* 2002; STUMPFERL *et al.* 2004). On the other hand, it was also shown that a high level of AOX due to the constitutive overexpression of the alternative oxidase in long-lived respiration mutants results in shortening life span and in restoring senescence (LORIN *et al.* 2001). We show here that neither the overexpression of the alternative oxidase in the short-lived  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-1* strain nor the inactivation of the *aox* resident gene in the long-lived  $\Delta$ *oxa1* (*oxa1<sup>ts</sup>*) *rmp1-2* strain significantly modifies the life span of these strains. This result clearly establishes that there is no causal relationship between the AOX level and life span.

The physiological role of the alternative pathway is still puzzling. It is of course important under conditions in which the cytochrome pathway is impaired. However, expression of the alternative oxidase can rescue mitochondrial respiration only if complex I is functional since the branch point is located downstream of this complex. We describe here for the first time a mutant displaying a defect in complex I activity in *P. anserina*. This defect is associated with a defect in complex IV and with a strong induction of the AOX, which, however, does not seem to play a crucial role in the control of life span since inactivation of the *aox* gene has no apparent effect. The induction of the AOX in a complex I-deficient mutant is, *a priori*, surprising and could only be a general response to a mitochondrial dysfunction, independent of its “usefulness” (which of course is evident for loss-of-function mutants of the cytochrome pathway). Although it was reported that ROS could in-

duce the expression of the alternative oxidase in plants (WAGNER 1995; YUKIOKA *et al.* 1998), the precise nature of the signal(s) that leads to this induction is unknown to date.

We previously showed that inactivation of complex IV in *P. anserina* results in an increase of life span associated with a decrease in the levels of free radicals (DUFOUR *et al.* 2000). Interestingly, the data obtained in this study also show that in strains mutated for OXA1, life span increase is correlated with ROS decrease. An impaired respiratory chain often increases oxidative stress (ISHII *et al.* 1998; McLENNAN and DEGLI ESPOSTI 2000; HARMAN 2001). The reverse effect observed in the  $\Delta\text{cox5}$  mutant of *P. anserina* was attributed to the presence of an alternative oxidase that could play a protective role in mitochondria by preventing ROS production. However, to date, this hypothesis has not been clearly demonstrated in *P. anserina* (LORIN *et al.* 2001). The reason(s) why deficiency in complex I in the  $\Delta\text{oxa1}$  ( $\text{oxa1}^{\Delta}$ ) *rmp1*-2 strain is associated with a reduction in ROS level is not yet well understood. Importantly, the presented data confirm the correlation between life-span increase and ROS decrease, supporting the idea that ROS level is a leading cause of aging.

Future studies will focus on deciphering the nature of the links between ROS production, AOX induction, and control of life span. They will also focus on the characterization of the interaction between the genes *oxa1* and *rmp1*. This characterization may hold a key for understanding the role of RMP1 in the control of mtDNA stability and for identification of its potential functional homologs in other species.

We thank M. Picard and L. Sperling for helpful discussions and comments on the manuscript. We gratefully acknowledge M. Minet and V. Rincheval for their help in the library construction and flow cytometric analysis, respectively. We also warmly thank V. Contamine for the gift of pHSS plasmid and T. Elthon and J. Velours for their gift of antibodies against AOX and  $\beta$ -subunit of ATPase, respectively. We are grateful to C. Barreau for providing chromoslot membranes and to M. Kermorgant and A. Léon for their help in the plasmid construction. This work was supported by grants from the Association Française contre les Myopathies.

#### LITERATURE CITED

- ALBERT, B., and C. H. SELLEM, 2002 Dynamics of the mitochondrial genome during *Podospora anserina* aging. *Curr. Genet.* **40**: 365–373.
- ALTAMURA, N., N. CAPITANIO, N. BONNEFOY, S. PAPA and G. DUJARDIN, 1996 The *Saccharomyces cerevisiae* OXA1 gene is required for the correct assembly of cytochrome c oxidase and oligomycin-sensitive ATP synthase. *FEBS Lett.* **382**: 111–115.
- BARREAU, C., C. SELLEM, P. SILAR, A. SAINARD-CHANET and B. TURCQ, 2002 A rapid and efficient method using chromoslots to assign any newly cloned DNA sequence to its cognate chromosome in the filamentous fungus *Podospora anserina*. *FEMS Microbiol. Lett.* **216**: 55–60.
- BAUER, M., M. BEHRENS, K. ESSER, G. MICHAELIS and E. PRATJE, 1994 PET1402, a nuclear gene required for proteolytic processing of cytochrome oxidase subunit 2 in yeast. *Mol. Gen. Genet.* **245**: 272–278.
- BELCOUR, L., O. BEGEL and M. PICARD, 1991 A site-specific deletion in mitochondrial DNA of *Podospora* is under the control of nuclear genes. *Proc. Natl. Acad. Sci. USA* **88**: 3579–3583.
- BELCOUR, L., A. SAINARD-CHANET, C. JAMET-VIERNY and M. PICARD, 1999 Stability of the mitochondrial genome of *Podospora anserina* and its genetic control, pp. 209–228 in *Mitochondrial Disease*, edited by P. LESTIENNE. Springer-Verlag, Berlin/New York.
- BERGES, T., and C. BARREAU, 1989 Heat shock at an elevated temperature improves transformation efficiency of protoplasts from *Podospora anserina*. *J. Gen. Microbiol.* **135**: 601–604.
- BONNEFOY, N., F. CHALVET, P. HAMEL, P. P. SLONIMSKI and G. DUJARDIN, 1994a OXA1, a *Saccharomyces cerevisiae* nuclear gene whose sequence is conserved from prokaryotes to eukaryotes controls cytochrome oxidase biogenesis. *J. Mol. Biol.* **239**: 201–212.
- BONNEFOY, N., M. KERMORGANT, O. GROUDINSKY, M. MINET, P. P. SLONIMSKI *et al.*, 1994b Cloning of a human gene involved in cytochrome oxidase assembly by functional complementation of an *oxa1*-mutation in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **91**: 11978–11982.
- BONNEFOY, N., M. KERMORGANT, O. GROUDINSKY and G. DUJARDIN, 2000 The respiratory gene OXA1 has two fission yeast orthologues which together encode a function essential for cellular viability. *Mol. Microbiol.* **35**: 1135–1145.
- BORGHOUTS, C., A. WERNER, T. ELTHON and H. D. OSIEWACZ, 2001 Copper-modulated gene expression and senescence in the filamentous fungus *Podospora anserina*. *Mol. Cell. Biol.* **21**: 390–399.
- BORGHOUTS, C., C. Q. SCHECKHUBER, A. WERNER and H. D. OSIEWACZ, 2002 Respiration, copper availability and SOD activity in *P. anserina* strains with different life span. *Biogerontology* **3**: 143–153.
- BRYAN, A. C., M. S. RODEHEFFER, C. M. WEARN and G. S. SHADEL, 2002 Sls1p is a membrane-bound regulator of transcription-coupled processes involved in *Saccharomyces cerevisiae* mitochondrial gene expression. *Genetics* **160**: 75–82.
- CONTAMINE, V., G. LECILLIER, L. BELCOUR and M. PICARD, 1996 Premature death in *Podospora anserina*: sporadic accumulation of the deleted mitochondrial genome, translational parameters and innocuity of the mating types. *Genetics* **144**: 541–555.
- CONTAMINE, V., D. ZICKLER and M. PICARD, 2004 The *Podospora rmp1* gene implicated in nucleus-mitochondria cross-talk encodes an essential protein whose subcellular location is developmentally regulated. *Genetics* **166**: 135–150.
- COPPIN, E., and R. DEBUCHY, 2000 Co-expression of the mating-type genes involved in internuclear recognition is lethal in *Podospora anserina*. *Genetics* **155**: 657–669.
- DEBUCHY, R., S. ARNAISE and G. LECILLIER, 1993 The mat-allele of *Podospora anserina* contains three regulatory genes required for the development of fertilized female organs. *Mol. Gen. Genet.* **241**: 667–673.
- D'ENFERT, C., M. MINET and F. LACROUTE, 1995 Cloning plant genes by complementation of yeast mutants. *Methods Cell Biol.* **49**: 417–430.
- DEQUARD-CHABLAT, M., and C. H. SELLEM, 1994 The S12 ribosomal protein of *Podospora anserina* belongs to the S19 bacterial family and controls the mitochondrial genome integrity through cytoplasmic translation. *J. Biol. Chem.* **269**: 14951–14956.
- DEQUARD-CHABLAT, M., E. COPPIN-RAYNAL, M. PICARD-BENNOUN and J. J. MADJAR, 1986 At least seven ribosomal proteins are involved in the control of translational accuracy in a eukaryotic organism. *J. Mol. Biol.* **190**: 167–175.
- DILLIN, A., A. L. HSU, N. ARANTES-OLIVEIRA, J. LEHRER-GRAIWER, H. HSIN *et al.*, 2002 Rates of behavior and aging specified by mitochondrial function during development. *Science* **298**: 2398–2401.
- DUFOUR, E., J. BOULAY, V. RINCHEVAL and A. SAINARD-CHANET, 2000 A causal link between respiration and senescence in *Podospora anserina*. *Proc. Natl. Acad. Sci. USA* **97**: 4138–4143.
- ELTHON, T., R. L. NICKELS and L. MCINTOSH, 1989 Monoclonal antibodies to the alternative oxidase of higher plants mitochondria. *Plant Physiol.* **89**: 1311–1317.
- ESSER, K., 1974 *Podospora anserina*, pp. 531–551 in *Handbook of Genetics*, edited by R. C. KING, Plenum Press, New York.
- FENG, J., F. BUSSIERE and S. HEKIMI, 2001 Mitochondrial electron transport is a key determinant of life span in *Caenorhabditis elegans*. *Dev. Cell* **1**: 633–644.
- HAMEL, P., W. SAKAMOTO, H. WINTZ and G. DUJARDIN, 1997 Functional complementation of an *oxa1*- yeast mutation identifies an

- Arabidopsis thaliana* cDNA involved in the assembly of respiratory complexes. *Plant J.* **12**: 1319–1327.
- HAMEL, P., C. LEMAIRE, N. BONNEFOY, P. BRIVET-CHEVILLOTTE and G. DUJARDIN, 1998 Mutations in the membrane anchor of yeast cytochrome c1 compensate for the absence of Oxa1p and generate carbonate-extractable forms of cytochrome c1. *Genetics* **150**: 601–611.
- HARMAN, D., 1956 Aging: a theory based on free radical and radiation chemistry. *J. Gerontol.* **11**: 298–300.
- HARMAN, P., 2001 Mitochondrial mutations differentially affect aging, mutability and anesthetic sensitivity in *Caenorhabditis elegans*. *Mech. Ageing Dev.* **122**: 1187–1201.
- HERRMANN, J. M., and W. NEUPERT, 2003 Protein insertion into the inner membrane of mitochondria. *IUBMB Life* **55**: 219–225.
- ISHII, N., M. FUJII, P. S. HARTMAN, M. TSUDA, K. YASUDA *et al.*, 1998 A mutation in succinate dehydrogenase cytochrome b causes oxidative stress and ageing in nematodes. *Nature* **394**: 694–697.
- JIA, L., M. DIENHART, M. SCHRAMP, M. MCCAULEY, K. HELL *et al.*, 2003 Yeast Oxa1 interacts with mitochondrial ribosomes: the importance of the C-terminal region of Oxa1. *EMBO J.* **22**: 6438–6447.
- KUHN, A., R. STUART, R. HENRY and R. E. DALBEY, 2003 The Alb3/Oxa1/YidC protein family: Membrane-localized chaperones facilitating membrane protein insertion? *Trends Cell Biol.* **13**: 510–516.
- LEE, S. S., R. Y. LEE, A. G. FRASER, R. S. KAMATH, J. AHRINGER *et al.*, 2003 A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat. Genet.* **33**: 40–48.
- LORIN, S., E. DUFOUR, J. BOULAY, O. BEGEL, S. MARSY *et al.*, 2001 Overexpression of the alternative oxidase restores senescence and fertility in a long-lived respiration-deficient mutant of *Podospora anserina*. *Mol. Microbiol.* **42**: 1259–1267.
- MCLENNAN, H. R., and M. DEGLI ESPOSTI, 2000 The contribution of mitochondrial respiratory complexes to the production of reactive oxygen species. *J. Bioenerg. Biomembr.* **32**: 153–162.
- MEYER, W., M. BAUER and E. PRATJE, 1997a A mutation in cytochrome oxidase subunit 2 restores respiration of the mutant pet ts1402. *Curt. Genet.* **31**: 401–407.
- MEYER, W., U. BOMER and E. PRATJE, 1997b Mitochondrial inner membrane bound Pet1402 protein is rapidly imported into mitochondria and affects the integrity of the cytochrome oxidase and ubiquinol-cytochrome c oxidoreductase complexes. *Biol. Chem.* **378**: 1373–1379.
- MUNKRES, K., 1990 Histochemical detection of superoxide radicals and hydrogen peroxide by Age-1 mutants of *Neurospora*. *Fungal Genet. Newsl.* **37**: 24–25.
- NARGANG, F. E., M. PREUSS, W. NEUPERT and J. M. HERRMANN, 2002 The Oxa1 protein forms a homooligomeric complex and is an essential part of the mitochondrial export translocase in *Neurospora crassa*. *J. Biol. Chem.* **277**: 12846–12853.
- NIJTMANS, L. G., N. S. HENDERSON and I. J. HOLT, 2002 Blue Native electrophoresis to study mitochondrial and other protein complexes. *Methods* **26**: 327–334.
- RIZET, G., 1952 Les phénomènes de barrage chez *Podospora anserina*. I. Analyse génétique des barrages entre souches Set s. *Rev. Cytol. Biol. Vég.* **13**: 51–92.
- RIZET, G., 1953 Impossibility of obtaining uninterrupted and unlimited multiplication of the ascomycete *Podospora anserina*. *C. R. Hebd. Seances Acad. Sci.* **237**: 838–840.
- RIZET, G., and C. ENGELMANN, 1949 Contribution à l'étude génétique d'un ascomycète tetraspore: *Podospora*. *Rev. Cytol. Biol. Vég.* **11**: 201–304.
- RODEHEFFER, M. S., and G. S. SHADEL, 2003 Multiple interactions involving the amino-terminal domain of yeast mtRNA polymerase determine the efficiency of mitochondrial protein synthesis. *J. Biol. Chem.* **278**: 18695–18701.
- ROUILLARD, J. M., M. E. DUFOUR, B. THEUNISSEN, E. MANDART, G. DUJARDIN *et al.*, 1996 SLS1, a new *Saccharomyces cerevisiae* gene involved in mitochondrial metabolism, isolated as a synthetic lethal in association with an SSM4 deletion. *Mol. Gen. Genet.* **252**: 700–708.
- SCHAGGER, H., and G. VON JAGOW, 1991 Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Anal. Biochem.* **199**: 223–231.
- SELLEM, C. H., A. SAINARD-CHANET and L. BELCOUR, 1990 Detection of a protein encoded by a class II mitochondrial intron of *Podospora anserina*. *Mol. Gen. Genet.* **224**: 232–240.
- SHIGENAGA, M. K., T. M. HAGEN and B. N. AMES, 1994 Oxidative damage and mitochondrial decay in aging. *Proc. Natl. Acad. Sci. USA* **91**: 10771–10778.
- SILAR, P., 1995 Two new easy to use vectors for transformation. *Fungal Genet. Newsl.* **42**: 73.
- SLUSE, F. E., and W. JARMUSZKIEWICZ, 1998 Alternative oxidase in the branched mitochondrial respiratory network: an overview on structure, function, regulation, and role. *Braz. J. Med. Biol. Res.* **31**: 733–747.
- STUMPFER, S. W., O. STEPHAN and H. D. OSIEWACZ, 2004 Impact of a disruption of a pathway delivering copper to mitochondria on *Podospora anserina* metabolism and life span. *Eukaryot. Cell* **3**: 200–211.
- SZYRACH, G., M. OTT, N. BONNEFOY, W. NEUPERT and J. M. HERRMANN, 2003 Ribosome binding to the Oxa1 complex facilitates co-translational protein insertion in mitochondria. *EMBO J.* **22**: 6448–6457.
- TRIFUNOVIC, A., A. WREDENBERG, M. FALKENBERG, J. N. SPELBRINK, A. T. ROVIO *et al.*, 2004 Premature ageing in mice expressing defective mitochondrial DNA polymerase. *Nature* **429**: 417–423.
- TURCQ, B., 1989 Clonage direct de gènes par complémentation chez le champignon *Podospora anserina*: application à l'étude de gènes d'incompatibilité. Ph.D. Thesis, University of Bordeaux II, Bordeaux, France.
- VANLERBERGHE, G. C., and L. MCINTOSH, 1997 Alternative oxidase: from gene to function. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**: 703–734.
- WAGNER, A. M., 1995 A role for active oxygen species as second messengers in the induction of alternative oxidase gene expression in *Petunia hybrida* cells. *FEBS Lett.* **368**: 339–342.
- YUKIOKA, H., S. INAGAKI, R. TANAKA, K. KATO, N. MIKI *et al.*, 1998 Transcriptional activation of the alternative oxidase gene of the fungus *Magnaporthe grisea* by a respiratory-inhibiting fungicide and hydrogen peroxide. *Biochim. Biophys. Acta* **1442**: 161–169.

Communicating editor: P. J. PUKKILA

