

# The Early Developmental Gene *Semaphorin 5c* Contributes to Olfactory Behavior in Adult *Drosophila*

Stephanie M. Rollmann,<sup>\*,†,1</sup> Akihiko Yamamoto,<sup>\*,†</sup> Tim Goossens,<sup>†</sup> Liesbeth Zwarts,<sup>†</sup>  
Zsuzsanna Callaerts-Végh,<sup>§</sup> Patrick Callaerts,<sup>‡,\*\*</sup> Koenraad Norga,<sup>‡,††</sup>  
Trudy F. C. Mackay<sup>†,‡‡</sup> and Robert R. H. Anholt<sup>\*,†,‡‡,2</sup>

<sup>\*</sup>Department of Zoology, <sup>‡‡</sup>Department of Genetics and <sup>†</sup>W. M. Keck Center for Behavioral Biology, North Carolina State University, Raleigh, North Carolina 27695, <sup>‡</sup>Laboratory of Developmental Genetics, Center for Human Genetics, B-3000 Leuven, Belgium, <sup>§</sup>Zoological Institute, Department of Biology, B-3000 Leuven, Belgium, <sup>\*\*</sup>Laboratory of Biological Psychology, B-3000 Leuven, Belgium and <sup>††</sup>Children's Hospital, B-3000 Leuven, Belgium

Manuscript received December 15, 2006

Accepted for publication April 13, 2007

## ABSTRACT

Behaviors are complex traits influenced by multiple pleiotropic genes. Understanding the mechanisms that give rise to complex behaviors requires an understanding of how variation in transcriptional regulation shapes nervous system development and how variation in brain structure influences an organism's ability to respond to its environment. To begin to address this problem, we used olfactory behavior in *Drosophila melanogaster* as a model and showed that a hypomorphic transposon-mediated mutation of the early developmental gene *Semaphorin-5c* (*Sema-5c*) results in aberrant behavioral responses to the repellent odorant benzaldehyde. We fine mapped this effect to the *Sema-5c* locus using deficiency mapping, phenotypic reversion through *P*-element excision, and transgenic rescue. Morphometric analysis of this *Sema-5c* allele reveals subtle neuroanatomical changes in the brain with a reduction in the size of the ellipsoid body. High-density oligonucleotide expression microarrays identified 50 probe sets with altered transcriptional regulation in the *Sema-5c* background and quantitative complementation tests identified epistatic interactions between nine of these coregulated genes and the transposon-disrupted *Sema-5c* gene. Our results demonstrate how hypomorphic mutation of an early developmental gene results in genome-wide transcriptional consequences and alterations in brain structure accompanied by profound impairment of adult behavior.

**B**EHAVIORS are complex traits influenced by the expression of multiple pleiotropic genes (ANHOLT 2004; ANHOLT and MACKAY 2004). Understanding the mechanisms that give rise to complex behaviors requires an understanding of how variation in transcriptional regulation shapes nervous system development and how variation in brain structure influences the ability of an organism to respond appropriately to its environment. To begin to address this problem, we used olfactory behavior in *Drosophila melanogaster* as a model, since chemosensation is essential for survival, the olfactory system of *Drosophila* has been well characterized (VOSSHALL 2000; DAHANUKAR *et al.* 2005), and flies are readily amenable to powerful genetic, neuroanatomical, and behavioral analyses. Previously, we showed that the genetic architecture of odor-guided behavior in this system is composed of epistatic networks of pleiotropic genes (FEDOROWICZ *et al.* 1998;

ANHOLT *et al.* 2003), that such networks are dynamic, and that genes functioning early in development feature prominently in genetic ensembles that mediate adult olfactory behavior (SAMBANDAN *et al.* 2006). Here, we characterize the effects of hypomorphic disruption of the early developmental gene, *Semaphorin-5c* (*Sema-5c*), on the transcriptome, on development of the central nervous system, and on adult olfactory behavior.

Semaphorins are a family of secreted and membrane-associated proteins studied extensively for their role in nervous system development, especially in axon guidance (KOLODKIN *et al.* 1992, 1993; LUO *et al.* 1993; PUSCHEL *et al.* 1995). Semaphorins have also been implicated in the function of the immune system (SHI *et al.* 2000; TAKEGAHARA *et al.* 2005) and a variety of diseases, including cancer (WOODHOUSE *et al.* 2003; NEUFELD *et al.* 2005; BASILE *et al.* 2006), retinal degeneration (RICE *et al.* 2004; ABID *et al.* 2006), schizophrenia (EASTWOOD *et al.* 2003), and rheumatoid arthritis (MILLER *et al.* 2004). Members of the semaphorin gene family are characterized by a conserved semaphorin domain ~500 amino acids in length (GHERARDI *et al.* 2004; YAZDANI and TERMAN 2006). The family consists

<sup>1</sup>Present address: Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45221.

<sup>2</sup>Corresponding author: W. M. Keck Center for Behavioral Biology, Campus Box 7617, North Carolina State University, Raleigh, NC 27695-7617. E-mail: anholt@ncsu.edu

of eight classes differing in their sequence and primary structures. Class 1 and 2 semaphorins have been identified solely in invertebrates, classes 3–7 are present in vertebrates with class 5 also found in *Drosophila*, and class 5 semaphorins occur only in viruses (SEMAPHORIN NOMENCLATURE COMMITTEE 1999).

Of particular interest to olfactory behavior is the implication that class 3 soluble semaphorins and their receptors play a role in axon guidance in the mouse olfactory system (RENZI *et al.* 2000; SCHWARTING *et al.* 2000; WALZ *et al.* 2002; TANIGUCHI *et al.* 2003; CLOUTIER *et al.* 2004). A null mutation in *Sema3F* resulted in defasciculation of the vomeronasal nerve and rerouting of axons from vomeronasal sensory neurons into the main olfactory bulb. Accurate innervation of the main olfactory bulb was also dependent on *Sema3F* signaling (CLOUTIER *et al.* 2004).

Of relevance to human disease are the class 5 semaphorins, which contain the characteristic semaphorin domain as well as seven thrombospondin repeat elements and a transmembrane domain (SEMAPHORIN NOMENCLATURE COMMITTEE 1999). A human class 5 semaphorin (*Semaf*) has been implicated in the cri-du-chat syndrome, a rare congenital neurological disorder (SIMMONS *et al.* 1998). The cri-du-chat critical region has been mapped to human chromosome 5 in which the human *Semaf* locus encompasses 10% of the interval. In *Drosophila*, a single class 5 semaphorin, *Sema-5c*, has been identified (KHARE *et al.* 2000; BAHRI *et al.* 2001) and is ubiquitously expressed in stage 2 embryos with a striped pattern emerging at later stages. Late-stage expression was observed at muscle attachment sites, the midgut, and the dorsal vessel (KHARE *et al.* 2000; BAHRI *et al.* 2001). *Sema-5c* expression patterns have not been characterized in the adult. Homozygous mutants of *Sema-5c* are viable with no detectable defects in embryonic development (BAHRI *et al.* 2001). Furthermore, in a search for suppressors of the *lethal giant larvae l(2)gl* phenotype, which exhibits neoplastic growth of the larval brain and imaginal discs, a *P*-element insertion near *Sema-5c* disrupted tumorigenesis (WOODHOUSE *et al.* 2003). A well-defined role for *Sema-5c* in neural development, however, has not yet been documented.

Recently, a *P*-element insertional mutagenesis screen for mutations affecting *Drosophila* olfactory avoidance behavior identified four independent *P*-element insertion lines in two different genetic backgrounds in which the *P* element inserted near the *Sema-5c* locus. All four *P*-element insertion lines showed aberrant olfactory avoidance behavior, suggesting that *Sema-5c* may play a role in odor-guided behavior (SAMBANDAN *et al.* 2006). Here, we characterize the role of *Sema-5c* in olfactory behavior by identifying and characterizing one of these *P*-element insertions in the *Sema-5c* gene region. We show that a *P*-element insertion upstream of the *Sema-5c* locus results in smell-impaired behavioral responses. We fine mapped this effect to the *Sema-5c* locus using

deficiency mapping, phenotypic reversion through *P*-element excision, and transgenic rescue. We found that mutation of the *Sema-5c* gene results in subtle changes in brain morphology with a reduction in the size of the ellipsoid body. Furthermore, we provide insights into the function of *Sema-5c* in adults by characterizing the genomewide transcriptional effects of the *P*-element insertion at the *Sema-5c* locus. These experiments demonstrate how hypomorphic mutation of an early developmental gene results in genomewide transcriptional consequences and alterations in brain structure accompanied by profound impairment of adult behavior.

## MATERIALS AND METHODS

***Drosophila* stocks:** The BG01245 line was generated and donated by Hugo Bellen as part of the Berkeley *Drosophila* Gene Disruption Project and has a *p[GT1]*-element insertion at cytological position 68F in the isogenic Canton-S (B) background (LUKACSOVICH *et al.* 2001; BELLEN *et al.* 2004). The BG01245 revertant line was generated as described below. The E215 and E77 excision lines and a *UAS-Sema-5c* transgenic line were generously made available by Sami M. Bahri (BAHRI *et al.* 2001). The transgenic construct was mobilized and placed into the Canton-S (B) background, as described below. The following mutants used for quantitative complementation tests for epistasis were obtained from the Bloomington Stock Center (stock number in parentheses): *no mitochondrial derivative* (BL-10715), *eukaryotic release factor 1* (BL-11488; BL-10266), *Sec61beta* (BL-12339; BL-10376), *CG3168* (BL-12516), *CG2994* (BL-12657), *CG17259* (BL-12894), *CG8533* (BL-13060), *laminin B1* (BL-13957), *CG4607* (BL-14408), *laminin A* (BL-14568), *CG4769* (BL-14909), *CG7800* (BL-14958), *CG8545* (BL-14964), between *CG5579/CG33177/CG33178* and *lipid storage droplet 2* (BL-15119), *lipid storage droplet 2* (BL-12398), *CG15557-9* (BL-15221), *CG8386* (BL-13815), *heterogeneous nuclear ribonucleoprotein at 27C* (BL-10375), *protein phosphatase 2A regulatory B subunit* (BL-12974), *signal sequence receptor beta* (BL-12094), *CG1383* (BL-15240), and *homothorax* (BL-11670). All flies were reared on standard agar–cornmeal–molasses medium in vials maintained at 25° and under a 12-hr light/dark cycle.

**Benzaldehyde avoidance assay:** All behavioral assays were conducted as described previously (ANHOLT *et al.* 1996). Briefly, one replicate assay consisted of a single sex group of five individuals, 5–7 days post-eclosion, removed from their food source ~1–2 hr prior to the assay. Test vials were demarcated at 3 and 6 cm from the bottom of the vial. Benzaldehyde was introduced on a saturated cotton swab wedged between the cotton plug and the vial wall at the 6-cm mark. Flies were allowed to acclimate to the vial for 15 sec after the introduction of the odor source. The number of flies present in the distal compartment of the vial was recorded every 5 sec for 1 min. The “avoidance score” for the replicate is the average of the 10 counts. All behavioral assays were conducted in an environmental chamber at 25° and multiple replicate assays were conducted for each genotype. Dose-response curves were determined using concentrations of 0.03, 0.06, 0.1, 0.3, 0.6, 1, and 3% (v/v) benzaldehyde, with subsequent behavioral assays conducted at 0.3% (v/v) benzaldehyde. Benzaldehyde avoidance responses of Canton-S (B) and BG01245 at different benzaldehyde concentrations were analyzed by a two-way fixed-effects ANOVA according to the model  $y = \mu + L + S + (L \times S) + E$ , where *L* denotes line, *S* designates sex, and *E* the environmental variation within sex and genotype. Ten

replicates per sex and genotype were measured at each benzaldehyde concentration.

**Quantitative complementation test:** The *P*-element insertion line (BG01245) and its co-isogenic control [Canton-S (B)] were crossed to excision lines E215 and E77 (BAHRI *et al.* 2001). Progeny from each cross were assayed for avoidance behavior to 0.3% (v/v) benzaldehyde and significant differences between phenotypic values were assessed between excision lines crossed to BG01245 and Canton-S (B). Forty replicates per genotype and sex were assayed over three blocks. For each complementation test, the data were analyzed by a three-way fixed-effects ANOVA with the model  $y = \mu + G + S + B + (G \times S) + (G \times B) + (S \times B) + (G \times S \times B) + E$ , with genotype (*G*), block (*B*), and sex (*S*) as fixed effects, and *E* indicating environmental variance. Quantitative failure to complement was inferred if the genotype or genotype-by-sex interaction ( $G \times S$ ) terms were significant.

**Phenotypic reversion through *P*-element excision:** The *p[GT1]* construct was mobilized by crossing BG01245 females to *w;Cy/Sp;SbΔ2-3/TM6,Tb* males. Male offspring of the genotype *w;Cy/CS(B);BG01245/SbΔ2-3* were subsequently mated to *w;CS(B);H/TM3,Sb* females, and single male offspring (*w;CS(B);P-/H*) were crossed to *w;CS(B);H/TM3,Sb* females. Male and female *w;CS(B);P-/TM3,Sb* in which the *P* element has been excised were mated *inter se* to make a homozygous *P*-element excision line in the isogenic Canton-S (B) background. Precise excision of the construct was characterized by PCR amplification using primers flanking the original *P*-element insertion site. PCR products were subsequently sequenced using ABI big dye terminator cycle sequencing chemistry (Applied Biosystems, Foster City, CA). Sequences were analyzed using Vector NTI Suite 9 software (Informax, Frederick, MD). Avoidance responses to benzaldehyde of the precise *P*-element excision line, Canton-S (B) control, and the BG01245 *P*-element insertion line were measured. Ten replicates per sex and line were scored and data were analyzed by a two-way fixed-effects ANOVA according to the model  $y = \mu + L + S + (L \times S) + E$ , where *L* denotes line, *S* designates sex, and *E* environmental variance. Significant differences among lines were determined by post-hoc Tukey's test.

**Transgenic rescue:** The *UAS-Sema-5c* transgene was mobilized and inserted into an isogenic Canton-S (B) background by crossing *+/+;UAS-Sema-5c* males to *w;Cy/Sp;TM3,Sb/H* females. Male offspring of the genotype *w;Cy/+;TM3/UAS-Sema-5c* were subsequently mated to *w;CS(B);TM3/H* virgin females and *w;Cy/CS(B);UAS-Sema-5c/TM3* female offspring were crossed to *w;Cy/Sp;SbΔ2-3/TM6,Tb* males. Male progeny, *w;Cy/CS(B);UAS-Sema-5c/SbΔ2-3*, were then crossed to *w;Cy/Sp;TM3/H* virgin females. Single males of the genotype *w;UAS-Sema-5c/Cy;SbΔ2-3/H* were subsequently mated to *w;Cy/Sp;TM3/H* females. Male progeny of the genotype *w;Cy/UAS-Sema-5c;TM3/H* were crossed to *w;Cy/Sp;TM3/H* females and *w;Cy/UAS-Sema-5c;TM3/H* offspring mated *inter se*. Male offspring, *w;UAS-Sema-5c;TM3/H*, were crossed to *w;Cy/Sp;BG01245* virgin females. Finally, males and females of the genotype *w;UAS-Sema-5c/Cy;BG01245/TM3* were mated *inter se*. Olfactory avoidance behavior to benzaldehyde was tested contemporaneously for the BG01245 *P*-element insertion line ( $N = 30$  replicates/sex), Canton-S (B) control ( $N = 30$  replicates/sex), and transgenic rescue line *w;UAS-Sema-5c/CS(B);BG01245* ( $N = 20$  replicates/sex). Avoidance responses were analyzed by a two-way fixed-effects ANOVA according to the model  $y = \mu + L + S + (L \times S) + E$ , where *L* denotes line, *S* designates sex, and *E* environmental variance. Significant differences among lines were determined by post-hoc Tukey's test.

**Whole-mount immunohistochemistry:** Brains were dissected in ice-cold phosphate buffered saline (PBS) for up to 1 hr and collected in PBS in a microcentrifuge tube on ice. All

the following steps were done on a rotating platform. After removal of PBS, the brains were fixed in PBS with 3.7% formaldehyde for 15 min at room temperature followed with three 10-min washes in PBS. The tissues were then preincubated with PAXD (PBS + 5% bovine serum albumin, 0.3% Triton-X100, and 0.3% sodium deoxycholate) for 10 min. Subsequently, the tissues were incubated overnight at 4° with PAXD containing the primary antibody at the appropriate dilution. Tissues were then washed for 4–6 hr with several changes of PAXD at room temperature and incubated overnight at 4° with PAXD with secondary antibody at the appropriate dilution. Tissues were washed for at least 2 hr with several changes prior to mounting in Vectashield medium (Vector Laboratories, Burlingame, CA).

**Antibody:** The monoclonal antibody 1D4 (antifasciclin2) was obtained from the Developmental Studies Hybridoma Bank (under the auspices of the National Institute of Child Health and Human Development and maintained by the University of Iowa, Department of Biological Sciences, Iowa City, IA 52242) and used at a dilution of 1:100. Cy3 and FITC-coupled secondary antibodies (Jackson ImmunoResearch, Westgrove, PA) were used at 1:200 and 1:100 dilutions.

**Microscopy:** The antifasciclin2 antibody staining used for morphometric analysis was documented using an Olympus BX61 epifluorescence microscope equipped with a DP70 digital camera controlled with analySIS software.

**Morphometric analysis:** For morphometric analyses, images were sampled using the analySIS software and the DP70 digital camera. Relevant dimensions (length and diameter of  $\alpha$ - and  $\beta$ -lobes and radii of ellipsoid body; see Figure 2D) were measured on the computer screen using a normal ruler and subsequently converted to values (expressed as percentages) relative to the distance between the two mushroom body heels. Statistical analyses used two-way ANOVA with post-hoc Tukey's tests.

**Transcriptional profiles:** For each genotype and sex, total RNA was isolated from two independent replicate groups of *Drosophila* heads (5–7 days post-eclosion) using Trizol (GIBCO-BRL, Gaithersburg, MD). RNA samples were purified through RNeasy columns (QIAGEN, Valencia, CA) and cDNA was generated from 5  $\mu$ g of total RNA. Biotinylated cRNA probes were generated and hybridized to *Drosophila* high-density oligonucleotide microarrays (GeneChip *Drosophila* Genome Array based on FlyBase version 1.0, Affymetrix) and visualized with a streptavidin–phycoerythrin conjugate according to Affymetrix protocols. The data were analyzed with Microarray Suite version 5.0 (MAS 5.0) using Affymetrix default analysis settings and global scaling as the normalization method. The signal values of any hybridization were multiplied by a scaling factor to make their mean intensity equal to 500. Differences in signal values between BG01245 and its control Canton-S (B) were analyzed by two-way factorial ANOVA according to the model  $y = \mu + L + S + (L \times S) + E$ , where *L* denotes line, *S* designates sex, and *E* the variance between replicate arrays. A significance threshold of  $P < 0.001$  was used to identify probe sets that differed between BG01245 and the Canton-S (B) control. Analysis of overrepresentation of molecular function and gene ontology categories among transcripts with altered transcriptional regulation was implemented with the DAVID program (DENNIS *et al.* 2003).

**Epistasis:** The *Sema-5c* *P*-element mutation is recessive. To assess epistatic interactions between genes with altered transcriptional regulation in the mutant background, *trans*-heterozygous flies were generated by crossing mutant lines from the Bloomington Stock Center to BG01245 and Canton-S (B). Ten replicates of five flies/sex and genotype were assayed for avoidance behavior to 0.3 and 0.6% (v/v) benzaldehyde as described above. Epistatic interactions were determined by ANOVA according to the model  $y = \mu + G + S + C + (G \times S) +$

$(G \times C) + (S \times C) + (G \times S \times C) + E$ , where  $G$  (genotype), sex ( $S$ ), and benzaldehyde concentration ( $C$ ) are fixed main effects and  $E$  indicates environmental variance.

## RESULTS

**A *P*-element insertion near the *Sema-5c* gene results in aberrant olfactory behavior:** Previously, in a screen for genes influencing olfactory avoidance behavior in *Drosophila*, four independent  $p[GT1]$ -element insertion lines adjacent to the *Sema-5c* locus showed aberrant behavioral responses to benzaldehyde (SAMBANDAN *et al.* 2006). The insertion BG01245 resulted in significantly reduced transcription levels of *Sema-5c* in larval and pupal development, whereas in the adult no such differences were observed (SAMBANDAN *et al.* 2006). We conducted a comprehensive characterization of the effects of this *P*-element insertion line on olfactory avoidance behavior (Figure 1). The *P*-element insertion in the BG01245 mutant line inserted 197 bp upstream of the first exon of *Sema-5c* (Figure 1A; SAMBANDAN *et al.* 2006). *Sema-5c* contains 12 exons and is ~14 kb in length with two predicted alternatively spliced transcripts that differ in the number of thrombospondin type 1 repeats (KHARE *et al.* 2000; BAHRI *et al.* 2001). Two predicted translation initiation sites have been identified in the first and second exons (BAHRI *et al.* 2001). In the BG01245 line, insertion of the  $p[GT1]$ -element into an isogenic Canton-S (B) background resulted in reduced avoidance responses to the standard test odorant benzaldehyde (Figure 1B). Reduced behavioral responses were similar for both sexes; therefore, measurements for males and females were pooled (Figure 1B). Dose responses to benzaldehyde for the mutant line compared to the control showed a decrease in avoidance behavior with a shift of the dose-response profile toward higher stimulus concentrations in the BG01245 mutant at concentrations of 0.1–3% (v/v) benzaldehyde (Figure 1B). Subsequent comparisons of avoidance responses between control and mutant flies were conducted at an intermediate concentration of 0.3% (v/v) benzaldehyde, which is optimal for resolving behavioral differences between the lines.

**$p[GT1]$ -element disruption of *Sema-5c* accounts for the smell-impaired phenotype:** To demonstrate that the smell-impaired phenotype is due to the  $p[GT1]$ -element insertion, we fine mapped the phenotype by quantitative complementation testing (PASYUKOVA *et al.* 2000; FANARA *et al.* 2002) using excision lines with small deletions in the *Sema-5c* gene, derived from an independent EP insertion line (Figure 1A; BAHRI *et al.* 2001). Results from these experiments allowed us to uncover genomic regions that fail to complement the mutant phenotype. The E77 and E215 excision lines differ in the sizes of their deletions with the E215 deletion spanning a portion of the 5'-untranslated region and first three exons of *Sema-5c*, whereas the E77 deletion removed a smaller portion of the coding region (see BAHRI *et al.* 2001; Figure

1A). Olfactory avoidance behavior of E77/Canton-S (B) flies was not significantly different from E77/BG01245 flies, indicating complementation of the mutant phenotype (Figure 1C). However, the E215/BG01245 genotype showed smell-impaired responses relative to the E215/Canton-S (B) genotype (Figure 1C;  $F_{1,148} = 6.84$ ,  $P < 0.0098$ ). Quantitative failure of the E215 excision to complement the BG01245 *P*-element insertion directly maps the effects of the mutation to the nonoverlapping region of the E77 and E215 deletions of the *Sema-5c* gene.

To further verify that aberrant olfactory avoidance responses are due to the  $p[GT1]$ -element insertion rather than to an independent mutation, we mobilized the  $p[GT1]$  transposon to obtain a precise *P*-element excision. Precise excision was verified by DNA sequencing the region around the original *P*-element insertion site. We observed significant differences in avoidance response between the control and the mutant, but not between the precise revertant and the control (Figure 1D;  $F_{2,54} = 32.53$ ,  $P < 0.0001$ ). Thus, precise excision of the  $p[GT1]$  construct resulted in restoration of avoidance responses to wild-type levels (Figure 1D).

Conclusive evidence that mutation of the *Sema-5c* gene is causal for the reduction in olfactory avoidance behavior was obtained by introducing an *UAS-Sema-5c* transgene into the BG01245 mutant background. The  $p[GT1]$  construct in the *Sema-5c* gene contains a *GAL4* cassette (LUKACSOVICH *et al.* 2001) that can be driven by the *Sema-5c* promoter, allowing transgenes cloned behind a *UAS* promoter to be expressed in cells that normally express the disrupted gene. The Canton-S (B) control, the BG01245 mutant, and the transgenic rescue line *w;UAS-Sema-5c/CS(B);BG01245* were tested contemporaneously. As expected, avoidance responses of Canton-S (B) differed significantly from those of the mutant (Figure 1E;  $F_{2,154} = 9.52$ ,  $P < 0.0001$ ). However, introduction of the *UAS-Sema-5c* transgene into the BG01245 genetic background resulted in rescue of avoidance behavior to control levels with no significant difference between the transgenic line and the Canton-S (B) control (Figure 1E). Quantitative complementation testing with deletions, phenotypic reversion through *P*-element excision, and transgenic rescue of avoidance responses to benzaldehyde show that disruption of the *Sema-5c* gene caused the observed smell-impaired phenotype.

**Neuroanatomical consequences of disruption of the *Sema-5c* gene:** In third instar Canton-S (B) larvae, we observed expression of *Sema-5c* in scattered cells in the brain and ventral ganglion and a broad pattern of expression at the level of the thoracic ganglion and in the midline glia (supplemental Figure 1 at <http://www.genetics.org/supplemental/>). However, we were not able to detect expression of *Sema-5c* in adult brain by *in situ* hybridization.

It was previously suggested that hypomorphic mutations due to insertion of  $p[GT1]$  might lead to subtle





FIGURE 1.—Phenotypic characterization of disruption of the *Sema-5c* gene by *p[GT1]* insertion. (A) *P*-element insertion site and deletions in the *Sema-5c* gene region. The horizontal line represents genomic DNA on the third chromosome at cytological position 68F2. Exons of the *Sema-5c* gene are represented by solid boxes. The location of the *p[GT1]* insertion site is indicated with an arrowhead and putative translation initiation sites are denoted by lines ending in circles. Excision lines E215 and E77 are represented as horizontal bars denoting the extent of the deletions of the *Sema-5c* gene region in these lines (BAHRI *et al.* 2001). (B) Dependence of olfactory avoidance behavior on the concentration of benzaldehyde. *P*-element insertion line BG01245 (shaded bars) and its control, Canton S (B) (solid bars), were measured at different benzaldehyde concentrations. The dashed red line denotes the expected baseline avoidance behavior score at which flies are indifferent to the odorant. Double asterisks denote  $P < 0.01$  at different concentrations of benzaldehyde (v/v) (0.1%  $F_{1,36} = 10.42$ ,  $P < 0.0027$ ; 0.3%  $F_{1,36} = 6.93$ ,  $P < 0.0124$ ; 0.6%  $F_{1,36} = 6.82$ ,  $P < 0.0131$ ; 1%  $F_{1,36} = 11.39$ ,  $P < 0.0018$ ; 3%  $F_{1,36} = 10.10$ ,  $P < 0.003$ ). (C) Fine mapping of aberrant olfactory avoidance responses in the BG01245 mutant to the *Sema-5c* gene region. Quantitative complementation tests were conducted with deletion lines E215 and E77 (A). Complementation is observed for deletion line E77. However, failure to complement is observed between progeny of crosses of Canton S (B) and BG01245 to E215, which spans both putative translation initiation sites of *Sema-5c*. The dashed red line denotes the expected baseline avoidance behavior score at which flies are indifferent to the odorant. No difference between the sexes was observed. Double asterisks denote  $P < 0.01$ . (D) Phenotypic reversion of avoidance responses to benzaldehyde by *P*-element excision. Precise excision of the *p[GT1]* construct restores the wild-type phenotype. Whereas *P*-element insertion line BG01245 shows reduced avoidance responses to 0.3% benzaldehyde compared with the control line Canton S (B), no significant difference is observed between Canton S (B) and the revertant line BG01245 Rev. Letters above bars denote results of the post-hoc Tukey's test with means designated by the same letter (a, b) not significantly statistically different from one another. The dashed red line denotes the expected baseline avoidance behavior score at which flies are indifferent to the odorant. (E) Transgenic rescue of the BG01245 aberrant behavioral phenotype. A full-length copy of the *UAS-Sema-5c* transgene, provided by Sami Bahri (BAHRI *et al.* 2001), was hooked to the second chromosome and placed into the co-isogenic Canton S (B) background (see MATERIALS AND METHODS). Canton S (B), BG01245, and the transgenic line *w;UAS-Sema-5c/CS(B);BG01245* were measured contemporaneously for avoidance behavior to 0.3% benzaldehyde. The dashed red line denotes the expected baseline avoidance behavior score at which flies are indifferent to the odorant. Letters above bars denote results of the post-hoc Tukey's test with different letters (a, b) denoting statistically significant mean line differences.



**FIGURE 2.**—Morphometric analysis of antifasciclin2 (1D4) antibody-stained adult brains of Canton S (B) control flies and BG01245 *Sema-5c* mutants. (A) Antifasciclin2 labels  $\alpha$ -,  $\beta$ -, and gamma ( $\gamma$ ) lobes as well as the ellipsoid body (eb) (B). (C) Short  $\alpha$ -lobe phenotype that was observed in 7 of 40 brains analyzed. Note that in the morphometric analysis brains were scored per scorable hemisphere (*i.e.*, without mechanical defects caused by dissections). Expressed per hemisphere, a total of 77 hemispheres were scorable, of which 7 displayed a readily identifiable short  $\alpha$ -lobe (arrow). This phenotype is never seen in wild-type flies. (D) Schematic of the adult *Drosophila* brain. For  $\alpha$ - and  $\beta$ -lobes, the lengths were measured at the positions indicated (double arrows marked with  $\alpha$  and  $\beta$ , respectively). The widths of the  $\alpha$ - and  $\beta$ -lobes were measured at the positions indicated by the black lines perpendicular to the lobes. For the ellipsoid body (in blue), two diameters were measured, EB-D1 and EB-D2, respectively. From these, the radii R1 and R2 were derived and the product  $R1 \times R2$  was used as a measure to compare surfaces of the ellipsoid body. All measurements were expressed as percentages relative to the distance between the heels (marked as the distance between  $\alpha$ -lobes). (E) No significant differences were observed between  $\alpha$ - or  $\beta$ -lobe diameters of Canton S(B) ( $N = 25$ , solid bars), *Sema-5c* (BG01245) mutants ( $N = 77$ , open bars), and *Sema-5c* revertants (BG01245 rev) ( $N = 32$ , shaded bars). (F) No significant differences were observed between  $\alpha$ - or  $\beta$ -lobe lengths of Canton S(B) ( $N = 25$ , solid bars), *Sema-5c* (BG01245) mutants ( $N = 77$ , open bars), and *Sema-5c* revertants (BG01245 rev) ( $N = 32$ , shaded bars). (G) A significant reduction ( $P < 0.01$ ) in ellipsoid body surface was found when control ( $N = 13$ , solid bar) and *Sema-5c* mutant (BG01245) ( $N = 40$ , open bar) ellipsoid body surfaces were compared. This phenotype is restored to wild type in the *Sema-5c* revertant (BG01245 rev) ( $N = 16$ , shaded bar).

morphological defects (SAMBANDAN *et al.* 2006). To examine whether disruption of the *Sema-5c* gene during development impacts the neuroanatomical organization of the adult brain, we performed whole-mount antibody labeling using the 1D4 monoclonal antibody against fasciclin 2 (*fas2*). Fasciclin 2 is strongly expressed in the ellipsoid body of the central complex, in the  $\alpha$ - and  $\beta$ -lobes of the mushroom bodies, and at somewhat lower levels in the gamma lobes of the mushroom bodies (Figure 2, A and B). This antibody staining allows good resolution for the identification of morphological defects in these neuropils. The gross anatomy of the *fas2*-positive neuropils was largely intact. We did, however, observe  $\alpha$ -lobe defects in  $\sim 10\%$  of the *Sema-5c* mutant brains (Figure 2C). These defects were never seen in the Canton-S (B) wild-type controls.

We measured the diameters and lengths of the  $\alpha$ - and  $\beta$ -lobes of the mushroom bodies and the surface of the ellipsoid body (Figure 2D). Morphometric analyses re-

vealed no significant differences in the diameters and lengths of the  $\alpha$ - and  $\beta$ -lobes of the mushroom bodies between controls, *Sema-5c* mutants (BG01245), and *Sema-5c* revertants (BG01245 rev) (Figure 2, E and F). We did, however, observe a significant reduction in surface area of the ellipsoid body, a phenotype that was completely reverted to wild type in the *P*-element excision revertant (BG01245 rev) (Figure 2G). A similar reversion of the phenotype was also seen in a transgenic rescue line, *w;UAS-Sema-5c/CS(B);BG01245* (results not shown). These experiments demonstrate that substantial impairments in adult behavior are associated with subtle alterations in brain structure due to disruption of the early developmental gene *Sema-5c*.

**Disruption of the *Sema-5c* gene alters genome-wide expression levels:** Previous studies have shown that the insertion of single *P* elements that affect olfactory avoidance behavior have genome-wide transcriptional consequences (ANHOLT *et al.* 2003). Therefore, we evaluated

to what extent disruption of the *Sema-5c* locus by the *P* element results in alterations in expression levels of transregulated genes. We assessed whole-genome transcriptional profiles of the BG01245 and Canton-S (B) lines by RNA hybridization to high-density oligonucleotide Affymetrix expression microarrays and used two-way ANOVA to partition variation in transcript abundance between main effects of sex and line (mutant *vs.* control genotypes) and the sex-by-line interaction term. We observed a total of 64 probe sets with significant difference in transcript abundance between genotypes and/or significant line-by-sex interaction terms at a significance threshold of  $P < 0.001$  (supplemental Table 1 at <http://www.genetics.org/supplemental/>; the conservative Bonferroni correction for multiple testing predicts that 14 of these coregulated genes may be false positives). A total of 43 probe sets were significant only for the line term, with 20 upregulated and 23 downregulated in the *Sema-5c* mutant background. Seven probe sets were significant for both line and sex-by-line interaction terms. For three of these probe sets, the direction of the effects was the same in males and females, and the interaction term was attributable to a change in relative magnitude of transcript between males and females. However, in four of these probe sets, and in the remaining 14 probe sets that were significant only for the sex-by-line interaction term, the changes in transcript abundance in the *Sema-5c* mutant were in opposite directions in males and females (supplemental Table 1 at <http://www.genetics.org/supplemental/>).

In these probe sets, genes that contribute directly to chemosensation included *antdh* (WANG *et al.* 1999), *homothorax* (CASARES and MANN 1998; DONG *et al.* 2000), *defective proboscis extension response 20* (NAKAMURA *et al.* 2002), and *odorant binding proteins* (*Obps*) *Obp56d*, *Obp99a*, *Obp99c*, and *Obp99d* (GALINDO and SMITH 2001; HEKMATSCAFE *et al.* 2002). Notably, the expression of *Obp99a* and *Obp99c* was downregulated in *Sema-5c* mutant males and upregulated in *Sema-5c* mutant females. Seven of the probe sets that differed in expression between lines represent predicted transcripts of unknown function. Gene ontology analysis identified overrepresented probe sets with altered transcript abundance in molecular function categories related to protein biosynthesis and metabolism, including purine nucleotide binding ( $P = 3.9E-2$ ), nucleotide binding ( $P = 7.4E-2$ ), oxidoreductase activity ( $P = 7.0E-2$ ), adenylyl nucleotide binding ( $P = 8.3E-2$ ), signal sequence binding ( $P = 4.7E-2$ ), elastase activity ( $P = 6.2E-2$ ), translation regulator activity ( $P = 6.7E-2$ ), translation factor activity/nucleic acid binding ( $P = 6.2E-2$ ), ATP binding ( $P = 7.0E-2$ ), as well as odorant binding ( $P = 1.4E-2$ ).

To investigate the extent to which genes with altered transcriptional regulation in the BG01245 mutant background interact epistatically with the transposon-tagged *Sema-5c* gene, we conducted quantitative complementa-

tion tests for epistasis (PASYUKOVA *et al.* 2000; FANARA *et al.* 2002; ANHOLT *et al.* 2003). We crossed the BG01245 mutant and its co-isogenic control to 24 available mutations (m) and tested the *trans*-heterozygous flies for avoidance behavior to 0.3 and 0.6% (v/v) benzaldehyde. We identified nine loci (37.5%) in which mutant alleles interacted epistatically with the *Sema-5c* *P*-element-tagged gene (Table 1). We found a significant reduction in avoidance behavior at 0.6% (v/v) benzaldehyde between mutants of coregulated genes crossed to the BG01245 mutant compared to the control for *heterogeneous nuclear ribonucleoprotein at 27C*, *lipid storage droplet 2*, *protein phosphatase 2A regulatory B subunit*, *CG1383*, and *homothorax*. *Eukaryotic release factor 1* and *Sec61beta* double heterozygotes were smell impaired at 0.3% (v/v) benzaldehyde relative to the control and double heterozygotes for *signal sequence receptor beta* showed reduced avoidance responses at both concentrations relative to the control genotype. Furthermore, the *CG8386* mutant showed a suppressor effect as a double heterozygote with the *Sema-5c* mutant when tested at 0.3% (v/v) benzaldehyde, but an enhancer effect at 0.6% (v/v) benzaldehyde (Table 1). Such plasticity of epistatic interactions and dependence on stimulus concentrations are consistent with previous observations (SAMBANDAN *et al.* 2006).

*Trans*-heterozygotes were also assayed for locomotor activity. Individual flies were placed into empty vials, given a gentle tap, and after acclimating for 15 sec were scored for locomotor activity over a 30-sec period. Nominally significant differences ( $P < 0.03$ ) were observed only for *trans*-heterozygotes of *CG8386* [ $27.96 \pm 0.34$  sec for m/Canton-S (B) and  $26.55 \pm 0.55$  sec for m/BG01245] and for *protein phosphatase 2A regulatory B subunit* [ $27.53 \pm 0.28$  sec for m/Canton-S (B) and  $26.24 \pm 0.51$  sec for m/BG01245]. No data were obtained for *CG1383* (BL-15240) and *homothorax* (BL-11670).

## DISCUSSION

We showed that *P*-element disruption of the early developmental gene *Sema-5c* results in smell-impaired behavioral responses to benzaldehyde. We mapped these behavioral effects to the *Sema-5c* gene using quantitative complementation tests to deletions, a precise *P*-element excision line, and transgenic rescue. *Sema-5c* mutants are homozygous viable and embryos do not show overt morphological defects (BAHRI *et al.* 2001).

To assess whether disruption of *Sema-5c* affects the structure of the adult brain, we performed morphometric measurements on the mushroom bodies and ellipsoid body, integrative centers in the central brain that could be readily visualized by staining with an antibody to fasciclin 2. We found neuroanatomical differences in the size of the ellipsoid body and in some cases the  $\alpha$ -lobes of the mushroom bodies. The observation that

TABLE 1

Significant epistatic interactions for olfactory avoidance behavior to benzaldehyde between BG01245 and loci with altered transcriptional regulation in the BG01245 mutant background

Gene name <sup>a</sup>	Benzaldehyde (% v/v)	Average avoidance score		$F_G$ statistic <sup>b</sup>	$P_G^c$	$P_{G \times C}^c$	$P_{G(0.3\%)}^d$	$P_{G(0.6\%)}^d$
		m × Canton S (B)	m × BG01245					
CG8386 (BL-13815)	0.3	2.77 ± 0.15;	3.19 ± 0.14;	0.06	0.8082	0.0046	0.0516	0.0403
	0.6	3.6 ± 0.15	3.11 ± 0.18					
Eukaryotic release factor 1 (BL-10266)	0.3	3.93 ± 0.14;	3.53 ± 0.14;	5.47	0.0222	0.4806	0.0311	0.2747
	0.6	4.29 ± 0.15	4.07 ± 0.14					
Heterogeneous nuclear ribonucleoprotein at 27C (BL-10375)	0.3	3.3 ± 0.16;	2.87 ± 0.22;	9.98	0.0023	0.5993	0.1159	0.0029
	0.6	3.78 ± 0.15	3.2 ± 0.11					
Lipid storage droplet 2 (BL-12398)	0.3	3.79 ± 0.18;	3.39 ± 0.22;	9.69	0.0027	0.303	0.1788	0.0031
	0.6	4.01 ± 0.13	3.21 ± 0.21					
Protein phosphatase 2A regulatory B subunit (BL-12974)	0.3	3.19 ± 0.13;	2.86 ± 0.13;	7.91	0.0063	0.6738	0.091	0.0322
	0.6	3.88 ± 0.14	3.44 ± 0.14					
sec61beta (BL-10376)	0.3	3.95 ± 0.11;	3.32 ± 0.17;	13.39	0.0005	0.1636	0.0036	0.0608
	0.6	4.2 ± 0.10	3.92 ± 0.10					
Signal sequence receptor beta (BL-12094)	0.3	3.56 ± 0.16;	2.75 ± 0.12;	24.25	<0.0001	0.5401	0.0004	0.0041
	0.6	3.76 ± 0.14	3.13 ± 0.14					
CG1383 (BL-15240)	0.3	3.04 ± 0.17;	2.84 ± 0.15;	5.34	0.0237	0.336	0.3814	0.0167
	0.6	3.7 ± 0.11	3.21 ± 0.16					
Homothorax (BL-11670)	0.3	3.6 ± 0.13;	3.39 ± 0.14;	5.1	0.027	0.3948	0.2713	0.0532
	0.6	3.7 ± 0.16	3.24 ± 0.14					

m, mutant background.

<sup>a</sup>Bloomington *Drosophila* stock numbers are in parentheses.

<sup>b</sup>F-statistic for genotype.

<sup>c</sup>P-values for genotype and genotype-by-concentration interactions.

<sup>d</sup>P-values for genotype when concentrations of benzaldehyde are analyzed separately. Lines BL-10715, 11488, 12339, 12516, 12657, 12894, 13060, 13957, 14408, 14568, 14909, 14958, 14964, 15119, and 15221 did not show epistatic interactions with BG01245.

*Sema-5c* is expressed in numerous structures in the third instar larval brain (supplemental Figure 1 at <http://www.genetics.org/supplemental/>) and that mutations in *Sema-5c* are associated with mild neuroanatomical defects (Figure 2) suggests that *Sema-5c* is required for normal *Drosophila* brain development and function. Intriguing in this regard is the demonstration that *Sema-5c* is linked to activation of the *Dpp/Mothers against Dpp* (*Mad*) signal transduction pathway (WOODHOUSE *et al.* 2003). This suggests a possible developmental mechanism by which disruption of *Sema-5c* may lead to the neuroanatomical and behavioral phenotypes that we describe here. The *Dpp* pathway is also important for synaptic development (SWEENEY and DAVIS 2002; RAWSON *et al.* 2003; DUDU *et al.* 2006). Thus, the observed neuroanatomical differences and the behavioral defects could in part be due to abnormal synapse structure and function. Whereas we were not able to detect expression of *Sema-5c* in adult brains by *in situ* hybridization, a previous study reported expression of *Sema-5c* in adult heads by real-time quantitative PCR (SAMBANDAN *et al.* 2006). Thus, expression of *Sema-5c* in the antennae cannot be excluded and would be consistent with the observed altered regulation of expression of the odorant-binding

proteins *Obp56d*, *Obp99a*, and *Obp99d* (supplemental Table 1 at <http://www.genetics.org/supplemental/>).

A low level or absence of *Sema-5c* expression in the adult brain suggests that the altered neuroanatomical features in the central brain arise as a consequence of earlier developmental impairments. This would also be consistent with the observation by SAMBANDAN *et al.* (2006) that the biggest differences in *Sema-5c* expression levels between the control Canton-S(B) line and the *Sema-5c* transposon-induced mutation are observed in earlier developmental stages. Furthermore, the neuroanatomical defects that we documented are more likely glial than neuronal in origin since the pattern of expression of *Sema-5c* in larvae is consistent with expression in glia.

The genetic architecture of olfactory behavior in adult *Drosophila* depends on a dynamic epistatic genetic network (ANHOLT *et al.* 2003). Developmental genes, including *Sema-5c*, form part of this network (SAMBANDAN *et al.* 2006). To assess to what extent transposon-mediated disruption of *Sema-5c* affects the expression of coregulated genes, we performed transcriptional profiling. We found 50 probe sets with altered regulation in the *Sema-5c* mutant background (supplemental Table 1 at <http://www.genetics.org/supplemental/>). This is in

accordance with previous studies that have shown that the insertion of a single *P* element affecting olfactory avoidance behavior results in a similar number of genes with altered transcription (ANHOLT *et al.* 2003). Quantitative complementation tests revealed significant concordance between transcriptional coregulation and enhancer or suppressor effects on phenotypic values in *trans*-heterozygotes (Table 1), in line with previous observations (ANHOLT *et al.* 2003).

Gene ontology analysis of the altered transcriptional profile in the *Sema-5c* mutant implicates genes involved in protein biosynthesis, transport, and secretion. The genes listed in supplemental Table 1 at <http://www.genetics.org/supplemental/> can be organized in a cellular pathway that directs protein synthesis and secretion of, among others, odorant-binding proteins, in line with a possible role of *Sema-5c* in support cells of the adult antenna. It is important, however, to note that the observed transcriptional profile is not necessarily due to the effect of *Sema-5c* disruption on a homogeneous population of cells, but may arise from a heterogeneous spatial and temporal pattern of expression. In addition, although changes in the size of the ellipsoid body and occasionally in the structure of the mushroom bodies accompany observed effects on olfactory behavior, we cannot be certain that aberrant olfactory behavior in *Sema-5c* flies is entirely due to effects of the mutation on the central brain. The implication of the ellipsoid body in altered olfactory behavior suggests the intriguing possibility that the ellipsoid body might play a direct role in processing olfactory information. On the other hand, the association of alterations in the ellipsoid body with altered olfactory behavior may reflect the previously demonstrated role of the ellipsoid body and other central complex structures in the control of locomotion (MARTIN *et al.* 1999; STRAUSS 2002).

Our results demonstrate how hypomorphic disruption of an early developmental gene can alter adult brain structure and transcriptional networks that contribute to adult behavior.

We thank Sami Bahri for kindly providing *Drosophila* stocks and Chao-Qiang Lai (Tufts University, Boston) for hybridization of the Affymetrix microarray GeneChips. This work was supported by National Institutes of Health grants GM059469 (to R.R.H.A.) and GM045146 (to T.F.C.M.). K.N. is a Senior Clinical Investigator of the Foundation for Scientific Research-Flanders and L.Z. and T.G. are supported by the Flanders Interuniversity Institute for Biotechnology (VIB). This work was partially supported by a Federal Science Policy Return Grant (K.N.), by a Marie-Curie International Reintegration Grant (K.N.), and VIB. This is a publication of the W. M. Keck Center for Behavioral Biology at North Carolina State University.

#### LITERATURE CITED

- ABID, A., M. ISMAIL, S. Q. MEHDI and S. KHALIQ, 2006 Identification of novel mutations in the SEMA4A gene associated with retinal degenerative diseases. *J. Med. Genet.* **43**: 378–381.
- ANHOLT, R. R. H., 2004 Genetic modules and networks for behavior: lessons from *Drosophila*. *BioEssays* **26**: 1299–1306.
- ANHOLT, R. R. H., and T. F. C. MACKAY, 2004 Quantitative genetic analyses of complex behaviours in *Drosophila*. *Nat. Rev. Genet.* **5**: 838–849.
- ANHOLT, R. R. H., R. F. LYMAN and T. F. C. MACKAY, 1996 Effects of single *P*-element insertions on olfactory behavior in *Drosophila melanogaster*. *Genetics* **143**: 293–301.
- ANHOLT, R. R. H., C. L. DILDA, S. CHANG, J. J. FANARA, N. H. KULKARNI *et al.*, 2003 The genetic architecture of odor-guided behavior in *Drosophila*: epistasis and the transcriptome. *Nat. Genet.* **35**: 180–184.
- BAHRI, S. M., W. CHIA and X. YANG, 2001 Characterization and mutant analysis of the *Drosophila sema 5c* gene. *Dev. Dyn.* **221**: 322–330.
- BASILE, J. R., R. M. CASTILHO, V. P. WILLIAMS and J. S. GUTKIND, 2006 Semaphorin 4D provides a link between axon guidance processes and tumor-induced angiogenesis. *Proc. Natl. Acad. Sci. USA* **103**: 9017–9022.
- BELLEN, H. J., R. W. LEVIS, G. LIAO, Y. HE, J. W. CARLSON *et al.*, 2004 The BDGP gene disruption project: single transposon insertions associated with 40% of *Drosophila* genes. *Genetics* **167**: 761–781.
- CASARES, F., and R. S. MANN, 1998 Control of antennal versus leg development in *Drosophila*. *Nature* **392**: 723–726.
- CLOUTIER, J. F., A. SAHAY, E. C. CHANG, M. TESSIER-LAVIGNE, C. DULAC *et al.*, 2004 Differential requirements for semaphorin 3F and Slit-1 in axonal targeting, fasciculation, and segregation of olfactory sensory neuron projections. *J. Neurosci.* **24**: 9087–9096.
- DAHANUKAR, A., E. A. HALLEM and J. R. CARLSON, 2005 Insect chemoreception. *Curr. Opin. Neurobiol.* **15**: 423–430.
- DENNIS, G., JR., B. J. SHERMAN, D. A. HOSACK, J. YANG, W. GAO *et al.*, 2003 DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol.* **4**: P3.
- DONG, P. D., J. CHU and G. PANGANIBAN, 2000 Coexpression of the homeobox genes *Distal-less* and *homothorax* determines *Drosophila* antennal identity. *Development* **127**: 209–216.
- DUDU, V., T. BITTIG, E. ENTCHIEV, A. KICHEVA, F. JULICHER *et al.*, 2006 Postsynaptic mad signaling at the *Drosophila* neuromuscular junction. *Curr. Biol.* **16**: 625–635.
- EASTWOOD, S. L., A. J. LAW, I. P. EVERALL and P. J. HARRISON, 2003 The axonal chemorepellant semaphorin 3A is increased in the cerebellum in schizophrenia and may contribute to its synaptic pathology. *Mol. Psychiatry* **8**: 148–155.
- FANARA, J. J., K. O. ROBINSON, S. M. ROLLMANN, R. R. H. ANHOLT and T. F. C. MACKAY, 2002 *Vanoso* is a candidate quantitative trait gene for *Drosophila* olfactory behavior. *Genetics* **162**: 1321–1328.
- FEDOROWICZ, G. M., J. D. FRY, R. R. H. ANHOLT and T. F. C. MACKAY, 1998 Epistatic interactions between smell-impaired loci in *Drosophila melanogaster*. *Genetics* **148**: 1885–1891.
- GALINDO, K., and D. P. SMITH, 2001 A large family of divergent *Drosophila* odorant-binding proteins expressed in gustatory and olfactory sensilla. *Genetics* **159**: 1059–1072.
- GHERARDI, E., C. A. LOVE, R. M. ESNOUF and E. Y. JONES, 2004 The sema domain. *Curr. Opin. Struct. Biol.* **14**: 669–678.
- HEKMAT-SCAFE, D. S., C. R. SCAFE, A. J. MCKINNEY and M. A. TANOUYE, 2002 Genome-wide analysis of the odorant-binding protein gene family in *Drosophila melanogaster*. *Genome Res.* **12**: 1357–1369.
- KHARE, N., N. FASCETTI, S. DAROCHA, R. CHIQUET-EHRISMANN and S. BAUMGARTNER, 2000 Expression patterns of two new members of the semaphorin family in *Drosophila* suggest early functions during embryogenesis. *Mech. Dev.* **91**: 393–397.
- KOLODKIN, A. L., D. J. MATTHES, T. P. O'CONNOR, N. H. PATEL, A. ADMON *et al.*, 1992 Fasciclin IV: sequence, expression, and function during growth cone guidance in the grasshopper embryo. *Neuron* **9**: 831–845.
- KOLODKIN, A. L., D. J. MATTHES and C. S. GOODMAN, 1993 The semaphorin genes encode a family of transmembrane and secreted growth cone guidance molecules. *Cell* **75**: 1389–1399.
- LUKACSOVICH, T., Z. ASZTALOS, W. AWANO, K. BABA, S. KONDO *et al.*, 2001 Dual-tagging gene trap of novel genes in *Drosophila melanogaster*. *Genetics* **157**: 727–742.
- LUO, Y., D. RAIBLE and J. A. RAPER, 1993 Collapsin: a protein in the brain that induces the collapse and paralysis of neuronal growth cones. *Cell* **75**: 217–227.
- MARTIN, J. R., T. RAABE and M. HEISENBERG, 1999 Central complex substructures are required for the maintenance of locomotor activity in *Drosophila melanogaster*. *J. Comp. Physiol. A* **185**: 277–288.
- MILLER, L. E., C. WEIDLER, W. FALK, P. ANGELE, J. SCHAUMBURGER *et al.*, 2004 Increased prevalence of semaphorin 3C, a repellent

- of sympathetic nerve fibers, in the synovial tissue of patients with rheumatoid arthritis. *Arthritis Rheum.* **50**: 1156–1163.
- NAKAMURA, M., D. BALDWIN, S. HANNAFORD, J. PALKA and C. MONTELL, 2002 Defective proboscis extension response (DPR), a member of the Ig superfamily required for the gustatory response to salt. *J. Neurosci.* **22**: 3463–3472.
- NEUFELD, G., N. SHRAGA-HELED, T. LANGE, N. GUTTMANN-RAVIV, Y. HERZOG *et al.*, 2005 Semaphorins in cancer. *Front. Biosci.* **10**: 751–760.
- PASYUKOVA, E. G., C. VIEIRA and T. F. C. MACKAY, 2000 Deficiency mapping of quantitative trait loci affecting longevity in *Drosophila melanogaster*. *Genetics* **156**: 1129–1146.
- PUSCHEL, A. W., R. H. ADAMS and H. BETZ, 1995 Murine semaphorin D/collapsin is a member of a diverse gene family and creates domains inhibitory for axonal extension. *Neuron* **14**: 941–948.
- RAWSON, J. M., M. LEE, E. L. KENNEDY and S. B. SELLECK, 2003 *Drosophila* neuromuscular synapse assembly and function require the TGF-beta type I receptor saxophone and the transcription factor Mad. *J. Neurobiol.* **55**: 134–150.
- RENZI, M. J., T. L. WEXLER and J. A. RAPER, 2000 Olfactory sensory axons expressing a dominant-negative semaphorin receptor enter the CNS early and overshoot their target. *Neuron* **28**: 437–447.
- RICE, D. S., W. HUANG, H. A. JONES, G. HANSEN, G.-L. YE *et al.*, 2004 Severe retinal degeneration associated with disruption of semaphorin 4A. *Invest. Ophthalmol. Vis. Sci.* **45**: 2767–2777.
- SAMBANDAN, D., A. YAMAMOTO, J. J. FANARA, T. F. C. MACKAY and R. R. H. ANHOLT, 2006 Dynamic genetic interactions determine odor-guided behavior in *Drosophila melanogaster*. *Genetics* **174**: 1349–1363.
- SCHWARTING, G. A., C. KOSTEK, N. AHMAD, C. DIBBLE, L. PAYS *et al.*, 2000 Semaphorin 3A is required for guidance of olfactory axons in mice. *J. Neurosci.* **20**: 7691–7697.
- SEMAPHORIN NOMENCLATURE COMMITTEE, 1999 Unified nomenclature for the semaphorins/collapsins. *Cell* **97**: 551–552.
- SHI, W., A. KUMANOGOH, C. WATANABE, J. UCHIDA, X. WANG *et al.*, 2000 The class IV semaphorin CD100 plays nonredundant roles in the immune system: defective B and T cell activation in CD100-deficient mice. *Immunity* **13**: 633–642.
- SIMMONS, A. D., A. W. PUSCHEL, J. D. MCPHERSON, J. OVERHAUSER and M. LOVETT, 1998 Molecular cloning and mapping of human semaphorin F from the Cri-du-chat candidate interval. *Biochem. Biophys. Res. Commun.* **242**: 685–691.
- STRAUSS, R., 2002 The central complex and the genetic dissection of locomotor behaviour. *Curr. Opin. Neurobiol.* **12**: 633–638.
- SWEENEY, S. T., and G. W. DAVIS, 2002 Unrestricted synaptic growth in spinster, a late endosomal protein implicated in TGF-beta-mediated synaptic growth regulation. *Neuron* **36**: 403–416.
- TAKEGAHARA, N., A. KUMANOGOH and H. KIKUTANI, 2005 Semaphorins: a new class of immunoregulatory molecules. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **360**: 1673–1680.
- TANIGUCHI, M., H. NAGAO, Y. K. TAKAHASHI, M. YAMAGUCHI, S. MITSUI *et al.*, 2003 Distorted odor maps in the olfactory bulb of semaphorin 3A-deficient mice. *J. Neurosci.* **23**: 1390–1397.
- VOSSHALL, L. B., 2000 Olfaction in *Drosophila*. *Curr. Opin. Neurobiol.* **10**: 498–503.
- WALZ, A., I. RODRIGUEZ and P. MOMBARTS, 2002 Aberrant sensory innervation of the olfactory bulb in neuropilin-2 mutant mice. *J. Neurosci.* **22**: 4025–4035.
- WANG, Q., G. HASAN and C. W. PIKIELNY, 1999 Preferential expression of biotransformation enzymes in the olfactory organs of *Drosophila melanogaster*, the antennae. *J. Biol. Chem.* **274**: 10309–10315.
- WOODHOUSE, E. C., A. FISHER, R. W. BANDLE, B. BRYANT-GREENWOOD, L. CHARBONEAU *et al.*, 2003 *Drosophila* screening model for metastasis: *Semaphorin 5c* is required for 1(2)gl cancer phenotype. *Proc. Natl. Acad. Sci. USA* **100**: 11463–11468.
- YAZDANI, U., and J. R. TERMAN, 2006 The semaphorins. *Genome Biol.* **7**: 211.

Communicating editor: L. HARSHMAN