

Table S2. Mean F_{ST} values between pairs of populations.

	Kenyan Bantu	S. African Bantu	Biaka Pygmy	Hadza	Maasai	Mandenka	Mbuti Pygmy	Namibian San	≠Khomani Bushmen	Sandawe	Yoruba
S. African Bantu	0.008										
Biaka	0.039	0.032									
Hadza	0.100	0.105	0.120								
Maasai	0.016	0.025	0.053	0.085							
Mandenka	0.015	0.015	0.046	0.108	0.028						
Mbuti	0.073	0.067	0.056	0.148	0.078	0.081					
Namibian Bushmen	0.091	0.073	0.074	0.153	0.090	0.097	0.092				
≠Khomani Bushmen	0.073	0.055	0.064	0.133	0.071	0.080	0.081	0.018			
Sandawe	0.021	0.025	0.050	0.085	0.015	0.033	0.075	0.082	0.065		
Yoruba	0.009	0.008	0.040	0.106	0.026	0.009	0.076	0.093	0.077	0.030	
Tuscan	0.135	0.147	0.172	0.180	0.095	0.146	0.193	0.198	0.172	0.108	0.148

Calculated over all autosomal SNPs. All populations are included except for the HapMap YRI, since the HGDP Yorubans are examined instead.