

**File S2: Cas9 preparation protocol  
(Paix *et al.* 2015)**

Protocol updates will be posted on the Seydoux lab website:

<http://www.bs.jhmi.edu/MBG/SeydouxLab/>

**Purification of Cas9::NLS<sub>SV40</sub>::His<sub>6</sub>:**

1. Transform DE3 GOLD (Agilent, #230132) cells with nm2973 plasmid (Fu *et al.* 2014) and plate on LB + 50 µg/mL Carbenicillin.
2. Inoculate 25mL LB + 50 µg/mL Carbenicillin with bacteria from the fresh transformation and incubate at 37°C overnight.
3. Transfer 5mL of overnight culture to 1L LB + 0.1% glucose + 50 µg/mL Carbenicillin and grow at 25°C. Grow to OD<sub>600</sub>~0.5.
3. Shift culture to 18°C for 15-25 minutes, then add IPTG to 0.2 mM. Incubate overnight.
4. Pellet culture and obtain wet weight. Resuspend at ~6 mL/g cells with Buffer A + protease inhibitor (Roche, #11836170001) + 1mM PMSF.
5. Sonicate 6 x 45s (setting 3 at 30%, 1 second pulse-2 second pause) with 1 minute cooling in between.
6. Spin lysate 30 minutes at 16000xg and transfer supernatant to a fresh tube.
7. Equilibrate a 5mL Ni-agarose (Qiagen, #30410) with column with Buffer A (at least 25mL).
8. Batch bind clarified lysate with Ni-agarose 45 minutes at 4°C.
9. Wash Ni-agarose column with 100mL of Buffer B.
10. Elute protein with Buffer C. Determine fractions that have Cas9 protein using Bradford assay or by running a small amount on SDS-PAGE gel. Pool fractions.
11. To remove contaminating DNA in the prep. Equilibrate a 5mL Q Sepharose (Sigma, #Q1126) column with 1M KCl (25mL, this charges the column). Then equilibrate Q Sepharose column with Buffer C (25mL).
12. Flow eluent (from step 11) over Q Sepharose column. Collect flow-through and dialyze into 1L Buffer D for 5 hours at 4°C. Transfer into 1L Buffer D and dialyze overnight.

13. Concentrate protein to ~10 mg/mL using a 100K centrifugal filter (Milipore, UFC910024). Aliquot and flash-freeze in liquid nitrogen. Store aliquots at -80°C. Typical yield is sufficient for 50-70 single-use aliquots (5µl aliquot, 10µg/µl Cas9).

**Buffers:**


Buffer A: 20mM Tris ph 8.0, 250 mM KCl, 20 mM imidazole, 10% glycerol, 1 mM TCEP

Buffer B: 20mM Tris ph 8.0, 800 mM KCl, 20 mM imidazole, 10% glycerol, 1mM TCEP

Buffer C: 20mM Hepes ph 8.0, 500 mM KCl, 250 mM imidazole, 10% glycerol

Buffer D: 20mM Hepes ph 8.0, 500 mM KCl, 20% glycerol

**Purified Cas9::NLS<sub>SV40</sub>::His<sub>6</sub> resolved by SDS-PAGE:**



Recombinant Cas9::NLS<sub>SV40</sub>::His<sub>6</sub> was affinity purified using Ni-agarose (lane 1). Pooled eluent was flowed over Q sepharose to remove contaminating DNA bound to Cas9 (lane 2). Samples were resolved by SDS-PAGE and visualized by coomassie staining.

**Cas9 activity assay:**

We recommend testing your Cas9 preparation using the method outlined in the direct delivery protocol (File S1, Section F).